

衛署藥製字第031991號

20'S

# "衛達" 優得舒 錠 10毫克

【成份】：每一錠含有：

Ritodrine Hydrochloride.....10mg

衛達化學製藥股份有限公司

408 台灣台中市工業區 23 路 21 號

【適應症】：  
預防早產、流產。

【注意】：  
請詳閱說明書後使用。

【用法用量】：

起始劑量1顆(10mg)約在靜脈注射停止前30分鐘給予。剛結  
藥的24小時內，每2個小時給一顆，之後，每4-6小時給1-2顆。  
這些劑量可根據子宮反應和副作用的出現而作調整。全天  
劑量不得超過120mg。治療期可隨醫師視需要而繼續給藥。  
本藥須由醫師處方使用。

製造批號: 127012  
生產日期: 20111219  
有效期至: 20161218



衛 達  
cGMP

DOH-PM-031991

20'S

# Utelax Tablets 10mg "Weidar"

【Composition】：Each tablet contains

Ritodrine Hydrochloride.....10mg

**WEIDAR CHEM. & PHARM. CO., LTD.**

N0.21, 23<sup>RD</sup> Road, Taichung Industrial Park, Taichung, Taiwan

【Indications】  
Prevention of premature labor or miscarriage.

【Dosage and Administration】  
Ritodrine hydrochloride be given by mouth in an initial dose of 10 mg may be given approximately 30 minutes before the termination of i.v. therapy. The usual dosage schedule for the first 24 hours of oral administration is 10 mg every 2 hours. Thereafter the usual maintenance dose is 10 to 20 mg every 4 to 6 hours, the dose depending on uterine activity and unwanted effects. The total daily dose of oral ritodrine should not exceed 120 mg. The treatment may be continued as long as the physician considers it desirable to prolong pregnancy.  
Use only on prescription by physician.



衛 達  
cGMP