

“國血製劑益康”

人類免疫球蛋白靜脈注射劑

“TBSF”

Human Immunoglobulin for Intravenous Use

本藥限由醫師使用

衛署簡效輸字第 000841 號

請與你的醫師討論使用此產品之風險與利益

藥商: 醫療財團法人台灣血液基金會

台北市南海路3號3樓

(02) 2351-1600

Store at 2°C to 8°C (Refrigerate. Do not freeze).
Protect from light.

The solution has a nominal osmolality of
375 mOsm/kg, is approximately isotonic and has
a pH of 4.25.

Inspect immediately before use. Do not use
if turbid.

Contains no antimicrobial preservative.
Use immediately on opening and discard any
unused portion appropriately.

Use in one patient on one occasion only.

Not for sale. “TBSF” Human Immunoglobulin for
Intravenous Use is manufactured from plasma
donated by Taiwan’s voluntary and
non-remunerated donors.

K37401850G/e/1

“TBSF”
Human Immunoglobulin
For Intravenous Use

3g in 50mL

(B)
EXP

“TBSF”
Human Immunoglobulin
For Intravenous Use

Solution for Intravenous Infusion

Human Immunoglobulin G 3g

3g in 50mL



CSL Behring

See package insert for Directions for Use.

This vial of “TBSF” Human Immunoglobulin for
Intravenous Use contains:

Human Immunoglobulin G	3g
(including protein Nitrogen	480 mg)
Human Immunoglobulin A	<1.25 mg
Maltose	14.6 mmol

Manufactured by CSL Behring (Australia) Pty Ltd
189-209 Camp Road Broadmeadows Vic 3047 Australia
Distributed by Taiwan Blood Services Foundation
3Fl. No.3 Nan-Hai Road Taipei 100 Taiwan R.O.C.