Action and Uses:
Effective gastric antacid of sustained action, with mild astringent and demulcent properties. For use in alleviating pain and hastening healing in duodenal or gastric ulcer.

Dosage:
Take 1 teaspoonful (10ml), 3–4 times daily. Please refer to the information contained inside for proper use.

SHAKE WELL BEFORE USE
KEEP CONTAINER TIGHTLY CLOSED

LOT:
EXP:

120 ML

複方 鋁膠懸液劑

ALU-GEL Suspension

GASTRIC ANTACID AGENT

Each ml contains:
Aluminum Hydroxide Gel, Dried.....50mg
Magnesium Trisilicate..... 100mg

M H

明華化學製藥廠股份有限公司
MING HWA DRUG CO., LTD.
台北縣瑞芳工業區頂坪路138號

【作　用】本品含有氫氧化鋁及三矽酸鎂兩種成分，能有效中和過多胃酸。單次劑量(10ml)之酸中和能力爲34.56mEq，鎂含量爲15.33mEq。

【適應症】緩解胃部不適或灼熱感、或經診斷爲胃及十二指腸潰瘍、胃炎、食道炎所伴隨之胃酸過多。

【用法・用量】
一天3至4次，飯後一小時及睡前或需要時服用。成人每次10ml，12歲以上適用成人劑量；6歲以上未滿12歲，每次5ml；3歲以上未滿6歲，每次2.5ml；3歲以下之嬰幼兒，請洽醫師診治，不宜自行使用。

請詳閱說明書
醫師藥師藥劑生指示藥品
內衛藥製字第09812號

使用前須充分搖勻