Agrylin®

(anagrelide HCl)

Platelet-reducing Agent

100 CAPSULES

Rx only

0.5 mg

Shire

DOSAGE AND ADMINISTRATION: The recommended starting dosage of AGRYLIN® is 0.5 mg bid or 2 capsules of 0.5 mg qd, which should be maintained for at least one week. Dosage may require adjustment. Total daily dose should not exceed 10 mg. See package outsert for complete information. Store at 25°C (77°F), in a light-resistant container, excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

000000

LOT 18432V7

EXP

批號

保存期限

Active Ingredients: Each capsule contains 0.5 mg of anagrelide base (as anagrelide hydrochloride).

Inactive Ingredients: anhydrous lactose, crospovidone, lactose monohydrate, magnesium stearate, microcrystalline cellulose and povidone.

安閱靈 0.5 毫克膠囊

衛署罕藥輸字第000007號 Agrylin 0.5mg Capsules

製造廠：Patheon Manufacturing Services LLC, 5900 Martin Luther King Jr.Highway, Greenville, NC 27834, USA

藥商：夏爾生技醫藥股份有限公司

地址：台北市信義區信義路五段7號80樓之1

本藥須由醫師處方使用

3 540092 063017