

Product Name: Aminomix Peripheral		Territory: EXP	Colour: ● Black ● Die cut	1. Draft 1. Corr	08.11.2016, 11:15 11.01.2017, 16:15		Variable Data:
Type of Packaging: Bag	Dosage: 2000 ml						
Material number: MOETxxxx/00 EXP	2-D-Matrix Code -						
Pharma-Code (Laetus) -	EAN Code: -						
Dimension: 280 x 370 mm	Font: Smallest Size: 9	Arial					
Operator: Roberto Grill +43(0) 699-17181288							



Aminomix Peripheral

Solution for infusion

780 kcal

2000 ml

Infusion via a peripheral or central vein

A two chamber bag (2000 ml) contains:

- 1. Amino acid solution with electrolytes:** L-Alanine 9.80 g, L-Arginine 8.40 g, Glycine 7.70 g, L-Histidine 2.10 g, L-Isoleucine 3.50 g, L-Leucine 5.18 g, L-Lysine 4.62 g, L-Methionine 3.02 g, L-Phenylalanine 3.57 g, L-Proline 7.84 g, L-Serine 4.55 g, Taurine 0.70 g, L-Threonine 3.08 g, L-Tryptophan 1.40 g, L-Tyrosine 0.28 g, L-Valine 4.34 g, Calcium chloride 0.36 g, Sodium glycerophosphate 3.56 g, Magnesium sulphate 1.56 g, Potassium chloride 2.82 g, Sodium acetate 2.32 g.
- 2. Glucose 12.6 %:** Glucose 126 g.

Corresponding to:
Amino acids 70 g, Nitrogen 11.4 g, Carbohydrates (glucose) 126 g, Electrolytes: sodium 50 mmol, potassium 38 mmol, magnesium 6.2 mmol, calcium 3.2 mmol, phosphorus 16.4 mmol, sulphate 6.2 mmol, chloride 44 mmol.

Energy content total	approx. 780 kcal
Energy content non protein	approx. 500 kcal
Osmolality:	approx. 860 mOsm/kg water
Osmolarity:	approx. 770 mmol/l
pH:	5.6 approx.

and acetic acid (pH-adjustment), water for injections.

For intravenous use by infusion via a peripheral or central vein.
Read the package leaflet before use.

Keep out of the sight and reach of children.

Any unused solution should be discarded.

Do not store above 30 °C.
Do not freeze.
Store in the overpouch.
Keep the bag in the outer carton to protect from light.

1. Remove the overpouch and lay the bag onto a solid surface with the ports pointing away.
2. Roll the bag straight from the top (handle) down to the ports until the peel seal has opened.

After separation of the peelable seal, the bag should be inverted repeatedly to ensure a homogenous mixture.

Do not use if package is damaged. Use only if the amino acid and glucose solutions are clear without particles and colourless to slightly yellow.

The content of the two separate chambers has to be mixed before use, and before any additions are made via the additive port.

Additions of other products to the mixed solution should be made aseptically, using the additive port.



MOETxxxx/00 EXP