

**【成 分】**  
Simvastatin...40mg

**【適應症】**

高膽固醇血症、高三酸甘油酯血症、降低冠心病高危險群或冠心病患者的心血管事件發生率及冠心病致死率、患有異核質家族性高膽固醇血症的兒童病患。

**【用法用量】及【注意事項】**

請詳閱說明書後使用。

本藥須由醫師處方使用。

請於25°C以下、避光儲存

請在有效期間內使用

逾期恕不接受退換

衛署藥製字第 052479 號



PIC/SG.M.P.& 國際ISO9001認證&中日合作廠

井田國際醫藥廠股份有限公司  
CHIN TENG INTERNATIONAL PHARMACEUTICAL MANUFACTURE CORP.  
台中市大甲幼獅工業區幼獅路32號  
No.32, Youshi Rd., Youshi Industrial Park Taja Taichung City, Taiwan

1000Tab

可治脂®  
膜衣錠  
40毫克

KEZHI®

FILM COATED TABLETS

40mg

本廠通過 PIC/S GMP 規範  
(衛署藥廠證字第 (AP) 0428050 號)

**【INDICATION】**

Primary hypercholesterolemia, heterozygous familial hypercholesterolemia or combined (mixed) hyperlipidemia in patients unresponsive to diet & non-pharmacological measures; to reduce the incidence of clinical coronary disease HDL-C in patients with primary hypercholesterolemia, heterozygous familial hypercholesterolemia or mixed hyperlipidemia. Treatment of patients with hypertriglyceridemia & primary dysbetalipoproteinemia. Adjunct to diet & other non-dietary measures to reduce elevated total-C, LDL-C & apo B in homozygous familial hypercholesterolemia.

**【DOSAGE & WARNINGS】**

Please read the instructions before use.

EXP:

LOT:



4 717216 018243

**【成 分】**  
Simvastatin...40mg

**【適應症】**

高膽固醇血症、高三酸甘油酯血症、降低冠心病高危險群或冠心病患者的心血管事件發生率及冠心病致死率、患有異核質家族性高膽固醇血症的兒童病患。

**【用法用量】及【注意事項】**

請詳閱說明書後使用。

本藥須由醫師處方使用。

請於25°C以下、避光儲存

請在有效期間內使用

逾期恕不接受退換

衛署藥製字第 052479 號



PIC/SG.M.P.& 國際ISO9001認證&中日合作廠

井田國際醫藥廠股份有限公司  
CHIN TENG INTERNATIONAL PHARMACEUTICAL MANUFACTURE CORP.  
台中市大甲幼獅工業區幼獅路32號  
No.32, Youshi Rd., Youshi Industrial Park Taja Taichung City, Taiwan

1000Tab

可治脂®  
膜衣錠  
40毫克

KEZHI®

FILM COATED TABLETS

40mg

本廠通過 PIC/S GMP 規範  
(衛署藥廠證字第 (AP) 0428050 號)

**【INDICATION】**

Primary hypercholesterolemia, heterozygous familial hypercholesterolemia or combined (mixed) hyperlipidemia in patients unresponsive to diet & non-pharmacological measures; to reduce the incidence of clinical coronary disease HDL-C in patients with primary hypercholesterolemia, heterozygous familial hypercholesterolemia or mixed hyperlipidemia. Treatment of patients with hypertriglyceridemia & primary dysbetalipoproteinemia. Adjunct to diet & other non-dietary measures to reduce elevated total-C, LDL-C & apo B in homozygous familial hypercholesterolemia.

**【DOSAGE & WARNINGS】**

Please read the instructions before use.

EXP:

LOT:



4 717216 018243