

AWB3BF0F1A79

Baxter

OLICLINOMEL N4-550 E

Emulsion for Infusion

1500 ml

FORMULA CONTAINING ELECTROLYTES

Lipid Emulsion (10%)

Volume 300 ml
Refined soya oil and refined olive oil* 30

Other ingredients:
Purified egg lecithin (1.2
Glycerol (3.7
Sodium oleate (0.0
Sodium Hydroxide (qs, pH
Water for Injections qs volume.

- Mixture of refined olive oil (approximately 80%) and refined soya oil (approximately 20%)
- For Peripheral and Central Intravenous Administration
- Keep all medicines out of reach / sight of children
- Use as directed by a medical practitioner

[illegible]醫生指示使用

Manufactured by:
[illegible]/AF SA
8, Boulevard René Branquart 7860 Lessines BELGIUM

[illegible] Marketed in [illegible]
da by:
[illegible] (India) Pvt. Ltd.
II Floor, Tower C, Building No. 8, IT Cyber City, DLF Phase II [illegible]gaon, Haryana – 122002 India
[illegible]: +91-124-4085291
[illegible] Lic. No.: FF- 94 -12052

[illegible]ured by:
[illegible] Healthcare (Asia) Pte Ltd, [illegible]gapore
[illegible] 412P

Imported by:
Baxter Healthcare (Thailand) Co., Ltd. Bangkok
Thai Reg. No.

ยาอันตราย

Amino Acids (5.5%) solution with electrolytes

	600 ml
Amino Acids	33 g
Sodium Acetate	1.47 g
Sodium glycerophosphate $5H_2O$	3.22 g
Potassium Chloride	1.79 g
Magnesium Chloride $6H_2O$	0.67 g
Calcium Chloride $6H_2O$	32 mmol
[illegible]	24 mmol
[illegible]	3.3 mmol
[illegible] sulfate	11 mmol
[illegible] ate	46 mmol
[illegible] ride	44 mmol

[illegible] ingredients:
[illegible]ic Acid (qs, pH), Water for Injections qs volume.

[illegible]centration of amino acids after mixing (per bag)

[illegible] amino acids:	2.41 g
[illegible]	1.85 g
[illegible]ucine	1.32 g
[illegible]nylalanine	1.91 g
[illegible]ethionine	(2.39 g)
[illegible]ine	1.98 g
[illegible]nine HCl)	1.91 g
[illegible]ycine	1.39 g
[illegible]ine	0.59 g
[illegible]rosine	
[illegible]ptophan	
[illegible] essential amino acids:	6.83 g
[illegible]ine	3.80 g
[illegible]ine	3.40 g
[illegible]ine	1.58 g
[illegible]ine	2.24 g
[illegible]ine	1.65 g
[illegible]ine	0.13 g
[illegible]ine	

[illegible] use only in adults and in children more than 2 years of age.
[illegible] parenteral nutrition in adults and children above 2 years of age when oral or enteral [illegible]trition is impossible, inadequate or contra-indicated.
[illegible] use if the bag is undamaged, the non-permanent seals are intact and the amino acid and glucose solutions are clear. Do not use if separation of the emulsion is observed.
Do not connect in series.
[illegible] not administer before, simultaneously or after administration of blood through the same equipment.
[illegible] not exceed the recommended infusion rate.
[illegible] unused product or waste material and all necessary [illegible]ces must be discarded.
[illegible]rile, endotoxin free solution.
[illegible]sage and administration: see enclosed leaflet.

[illegible]piry date / ยาสิ้นอายุ

02-20[illegible]7

Glucose (20%) solution with calcium chloride

	600 ml
Volume	120 g
Anhydrous glucose	(132 g
(As glucose monohydrate)	0.44
Calcium Chloride	3 mmol
Ca^{++}	6 mmol
Chloride	

Other Ingredients:
Hydrochloric Acid (qs, pH)
Water for Injections qs volume.

After the contents of the three compartments have been mixed, the ternary mixture provides the following:

Volume	1500 ml
Osmolarity	750 mOsm/l
pH	[illegible]
Nitrogen	5.4
Amino acids	33
Total calories	910 kcal
Non protein calories	780 kcal
Glucose calories	480 kcal
Lipid calories	300 kcal
Non-protein calorie/nitrogen ratio	144 kcal/g
[illegible]	32 mmol
[illegible]	24 mmol
[illegible]	3.3 mmol
[illegible]	3 mmol
[illegible]	13 mmol
[illegible] phosphate**	46 mmol
[illegible]cetate	50 mmol
[illegible]oride	

[illegible] phosphates provided by the lipid emulsion.

[illegible]500 ml in 3 compartment bag; box of 4.
[illegible]ep container in the outer carton. Do not freeze.
[illegible]ly administer after breaking the non-permanent seals between the compartments. It is recommended that the reconstituted [illegible]mulsion is used immediately.
[illegible] reconstituted emulsion is stable for 7 days at [illegible]tween +2 °C and +8 °C followed by a maximum of [illegible] hours at temperatures not exceeding 25 °C.
[illegible]ure product is at room temperature.
[illegible]ditives may be incompatible. Refer to the manufacturer [illegible] further details.
[illegible]e ternary mixture is hypertonic and may cause vein [illegible]itation.

[illegible]er and OliClinomel are trademarks of [illegible]er International nc.

[illegible]ch N°

BE-35-01-820

15025N11

"百特" 歐諾美

N4-550E輸注乳液

衛署藥輸字第 024679 號
百特醫療產品股份有限公司
製造廠：Baxter S.A. 地址：BD.R. Branquart 80, 7860 Lessines, Belgium
封存層(藥水接觸層)材質為20%SEBS+80%EP，其他包材請詳如仿單
請貯存於 25°C 以下，不可冷凍
貼標廠:台灣大昌華嘉股份有限公司
地址:桃園縣楊梅市瑞坪里梅獅路二段629號

本藥限由醫師使用
台北市敦化南路二段216號15樓

包裝:1500ml
OLIN415CNL01

使用說明:去除外包裝，從軟袋頂端(吊孔端)開始用手將袋子捲起來，待封條撕開後，將軟袋上下顛倒讓內容物充分混合。
(操作方式詳見仿單)