仿單標籤粘貼表

產品名稱	欣保維優靜脈營養輸注液	申請廠商	台灣柏朗股份有限公司
衛生福利部食品藥物管理署給證號碼	衛署藥輸字 第 025553 號		

Nutriflex Omega special — **Emulsion for Infusion**

179/12612324/0819

欣保維優靜脈營養輸注液

1250 ml of the ready to use emulsion contain after mixing of the three chamber contents

from the upper left-hand chamber (500 ml):

Glucose monohydrate equivalent to anhydrous glucose 180.0 g	198.0 g
Sodium dihydrogen phosphate dihydrate	3.120 g
Zinc acetate dihydrate	8.78 mg

from the upper right-hand chamber (250 ml):

Medium-chain triglycerides	25.0 g
Soya-bean oil refined	20.0 g
Omega-3-acid triglycerides	5.0 g

from the lower chamber (500 ml):

Isoleucine	4.11 g
Leucine	5.48 g
Lysine hydrochloride equivalent to Lysine 3.98 g	4.98 g
Methionine	3.42 g
Phenylalanine	6.15 g
Threonine	3.18 g
Tryptophan	1.00 g
Valine	4.51 g
Arginine	4.73 g
Histidine hydrochloride monohydrate equivalent to Histidine 2.19 g	2.96 g
Alanine	8.49 g
Aspartic acid	2.63 g
Glutamic acid	6.14 g
Glycine (aminoacetic acid)	2.89 g
Proline	5.95 g
Serine	5.25 g
Sodium hydroxide	1.464 g
Sodium chloride	0.473 g
Sodium acetate trihydrate	0.313 g
Potassium acetate	4.611 g
Magnesium acetate tetrahydrate	1.137 g
Calcium chloride dihydrate	0.779 g

Electrolytes:

Sodium	67.0	mmol
Potassium	47.0	mmol
Magnesium	5.3	mmol
Calcium	5.3	mmol
Zinc	0.04	mmol
Chloride	60	mmol
Acetate	60	mmol
Phosphate	20	mmol

Amino acid content:	71.8 g
Total nitrogen content:	10.0 g
Carbohydrate content:	180.0 g
Lipid content:	50.0 g

Energy in the form of lipid 1990 kJ (475 kcal)

Energy in the form of carbohydrates 3015 kJ (720 kcal)

Energy in the form of amino acids 1170 kJ (280 kcal)

Non-protein energy 5005 kJ (1195 kcal)

Total energy 6175 kJ (1475 kcal)

Osmolality 2090 mOsm/kg

pH 5.0 – 6.0

Citric acid monohydrate, Egg lecithin, Glycerol, Sodium oleate, all-rac-α-Tocopherol, Sodium hydroxide for pH adjustment, Water for injections.

Intravenous use.

For central venous infusion only.

For single use.

Read the package leaflet before use.

Keep out of the reach and sight of children.

Only use bags that are undamaged and where is no discernible phase separation of the emulsion after gentle shaking and in which the amino acid and glucose solutions are clear.

After first opening the product should be used immediately.

Read the leaflet for the shelf life of the mixed product.

Do not store above 25°C.

Do not freeze. If accidentally frozen, discard the bag.

Keep the bags in the outer carton in order to protect from light.

衛署藥輸字第025553號

本藥限由醫師使用

NuTRIflex Omega special 使用具彈性且材質為polyamide/polypropylene之多腔室密封袋

4 030539 126700

製造廠：**B. Braun Melsungen AG**

廠　址：Carl-Braun-Straße 1
D-34212 Melsungen, Germany

公　司：D-34209 Melsungen, Germany

藥　商：台灣柏朗股份有限公司

地　址：台北市松山區健康路152號9樓

REF **354 0122**

GTIN: 04030539126700

5 x 1250 ml

Manuf. date:

LOT

EXP:

B. Braun Melsungen AG
34209 Melsungen
Germany

108. 11. 19

衛生福利部 食品藥物管理署 騎縫章

12612324_NuTRIflexOmega special_KET5x1250ml__TW_179.indd 1 02.08.19 11:01

※ 外文仿單應檢附中文譯文

（1）報核仿單標籤以粘貼全形實物為原則。

註

（2）仿單標籤等實物過大或印於玻璃金屬容器等不便於粘貼時附送現品並將照相影本代替粘貼報核


1250 ml of the ready to use emulsion contain after mixing of the three chamber contents
from this chamber (500 ml):

Glucose monohydrate	198.0 g
equivalent to anhydrous glucose 180.0 g	
Sodium dihydrogen phosphate dihydrate	3.120 g
Zinc acetate dihydrate	8.78 mg

欣保維優
靜脈營養輸注液
NuTRIflex
Omega special
Emulsion for Infusion

1250 ml of the ready to use emulsion contain after mixing of the three chamber contents
from this chamber (250 ml):

Medium-chain triglycerides	25.0 g
Soya-bean oil refined	20.0 g
Omega-3-acid triglycerides	5.0 g



4 030539 126656

Manuf. date: LOT EXP:

衛生福利部 食品藥物管理署 騎縫章

108. 11. 19

2000

欣保維優靜脈營養輸注液 Nutriflex Omega special
Emulsion for Infusion

1250 ml of the ready to use emulsion contain after mixing of the three chamber contents
from this chamber (500 ml):

Isoleucine	4.11 g
Leucine	5.48 g
Lysine hydrochloride	4.98 g
equivalent to Lysine 3.98 g	
Methionine	3.42 g
Phenylalanine	6.15 g
Threonine	3.18 g
Tryptophan	1.00 g
Valine	4.51 g
Arginine	4.73 g
Histidine hydrochloride monohydrate	2.96 g
equivalent to Histidine 2.19 g	
Alanine	8.49 g
Aspartic acid	2.63 g
Glutamic acid	6.14 g
Glycine (aminoacetic acid)	2.89 g
Proline	5.95 g
Serine	5.25 g
Sodium hydroxide	1.464 g
Sodium chloride	0.473 g
Sodium acetate trihydrate	0.313 g
Potassium acetate	4.611 g
Magnesium acetate tetrahydrate	1.137 g
Calcium chloride dihydrate	0.779 g

1500

1000

Electrolytes:	
Sodium	67.0 mmol
Potassium	47.0 mmol
Magnesium	5.3 mmol
Calcium	5.3 mmol
Zinc	0.04 mmol
Chloride	60 mmol
Acetate	60 mmol
Phosphate	20 mmol

Amino acid content:	71.8 g
Total nitrogen content:	10.0 g
Carbohydrate content:	180.0 g
Lipid content:	50.0 g

Energy in the form of lipid 1990 kJ (475 kcal)
Energy in the form of carbohydrates 3015 kJ (720 kcal)
Energy in the form of amino acids 1170 kJ (280 kcal)
Non-protein energy 5005 kJ (1195 kcal)
Total energy 6175 kJ (1475 kcal)

Osmolality	2090 mOsm/kg
pH	5.0 – 6.0

Citric acid monohydrate, Egg lecithin, Glycerol, Sodium oleate, all-rac-α-Tocopherol, Sodium hydroxide for pH adjustment, Water for injections.

Intravenous use.

For central venous infusion only.

For single use.

Read the package leaflet before use.

Keep out of the reach and sight of children.

Only use bags that are undamaged and where is no discernible phase separation of the emulsion after gentle shaking and in which the amino acid and glucose solutions are clear.

After first opening the product should be used immediately.

Read the leaflet for the shelf life of the mixed product.

Do not store above 25°C.
Do not freeze. If accidentally frozen, discard the bag.

Keep the bags in the outer carton in order to protect from light.

衛署藥輸字第
025553號

本藥限由醫師使用

NuTRIflex Omega special
使用具彈性且材質為
polyamide/polypropylene
之多腔室密封袋

1250 ml

500

B|BRAUN

B. Braun Melsungen AG
34209 Melsungen
Germany

製造廠 ： B. Braun Melsungen AG
廠　址 ： Carl-Braun-Straße 1,
D-34212 Melsungen, Germany
公　司 ： D-34209 Melsungen, Germany
藥　商 ： 台灣柏朗股份有限公司
地　址 ： 台北市松山區健康路152號9樓

179/FF65902/0819