

# Evaluation of Total Antioxidant Activity of Several Popular Vegetables and Chinese Herbs: A Fast Approach with ABTS/H<sub>2</sub>O<sub>2</sub>/HRP System in Microplates

ING-CHIEN CHEN<sup>1</sup>, HUI-CHI CHANG<sup>2</sup>, HUI-WEN YANG<sup>3</sup> AND GAN-LIN CHEN<sup>4\*</sup>

<sup>1</sup> Institute of Plant Biology, National Taiwan University, Taipei 106, Taiwan (R.O.C.)

<sup>2</sup> Division of Pharmacy, The Buddhist Tzu Chi General Hospital in Dalin, Chiayi 622, Taiwan (R.O.C.)

<sup>3</sup> Division of Pharmacy, Koo Foundation Sun Yat-Sen Cancer Center, Taipei 112, Taiwan (R.O.C.)

<sup>4</sup> Institute of Pharmaceutical Chemistry, China Medical University, Taichung 404, Taiwan (R.O.C.)

(Received: April 20, 2003; Accepted: September 26, 2003)

## ABSTRACT

Total antioxidant activities for several popular vegetables and traditional Chinese herbs were evaluated in an ABTS/H<sub>2</sub>O<sub>2</sub>/HRP system with a microplate reader. This improved method offers a rapid and sensitive way to measure total antioxidant activity of testing samples. Most samples had antioxidative capacities. Among the tested vegetables, sugar beet and red cabbage had the highest total antioxidant activity, whereas tropical Almond Terminalia and Moutan Radicis Cortex had exceptionally high total antioxidant activity in the tested Chinese herbs.

Key words: total antioxidant activity, vegetables, Chinese herb

## INTRODUCTION

Reactive oxygen species (ROS) are involved in the development of various diseases and also accelerate aging. ROS scavengers such as superoxide dismutase and antioxidants can effectively reduce damage from oxidation. Experimental evidence suggests that free radicals and ROS can take a part in many diseases<sup>(1,2)</sup>. As plants produce antioxidants to control the oxidative stress caused by sunlight and oxygen, they became a source of useful new compounds with antioxidant activity for human consumption. Traditional Chinese medicinal plants possessing natural antioxidants such as anthraquinones, flavonoids, aromatic acids, and tannins have recently been shown to have ROS scavenging and lipid peroxidation prevention effects<sup>(3-7)</sup>. Evidence is mounting for a role of dietary phytochemicals, including flavonoids, ascorbic acid,  $\alpha$ -tocopherol, and carotenoids, in the maintenance of health and protection from disease<sup>(8-14)</sup>. The epidemiological evidence for a protective effect of natural food and herb antioxidants against cancer, coronary heart disease, age-related macular degeneration, respiratory disease, cystic fibrosis, immuno disease, and Alzheimer's disease is impressive<sup>(15,16)</sup>. Numerous surveys have shown an inverse relationship between the intake of vegetables and the incidence of those disorders<sup>(17,18)</sup>.

The relationship between these discoveries and the mechanisms of traditional herbal therapies remains open to investigation. Chinese have used herbal medicine for over

four thousand years. Most herbs are boiled with water and the water extracts are then used for treatment. For example, the boiled soup of fallen Tropical Almond Terminalia Leaves is beneficial in treating hepatitis<sup>(19)</sup>. Moutan Radicis Cortex and Rehmanniae Radix et Rhizoma are claimed to remove heat from blood and activate blood circulation. Ligustri Lucidi Fructus, Asari Herba Radice, and Angelicae Dahuricae relieve pain by dispelling wind, coldness and dampness. Leonuri Herba is said to regulate menstruation by activating blood circulation and inducing diuresis<sup>(20)</sup>. The proportions of phytochemicals with antioxidative activity vary among foods. In order to investigate possible action mechanisms for such herbal remedies, a method to measure the total antioxidant activity in food and herb should be a substantial asset, regardless of the individual antioxidants which contribute toward this activity<sup>(21)</sup>.

Established methods for the measurement of total antioxidant activity are all essentially inhibition methods: a free radical species is generated, there is an end point by which the presence of the radical is detected, and the antioxidant activity of the added sample inhibits the end point by scavenging the free radical. The ABTS/H<sub>2</sub>O<sub>2</sub>/HRP system presented here is reliable and practical for measuring total antioxidant activity in fluid samples<sup>(22)</sup>. When ABTS (2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid)) is incubated with peroxidase and hydrogen peroxide, the relatively long-lived radical cation, ABTS<sup>+</sup>, is formed<sup>(23)</sup>. In the presence of antioxidant reductants and hydrogen donors, the radical cation is quenched and green color formation delayed. If the maximum ABTS<sup>+</sup> absorp-

\* Author for correspondence. Tel: 886-4-22053366 ext. 1914;  
Fax: 886-4-22031075; E-mail: glchen@mail.cmc.edu.tw

tion at 414 nm is traced, the lag time is proportional to the total antioxidant activity in the sample<sup>(21)</sup>. A microplate assay was reported<sup>(24)</sup> for determining superoxide dismutase. We had studied total antioxidant activity of plants for years, and in this report we intended to study that of several popular vegetables and traditional Chinese herbs in the ABTS/H<sub>2</sub>O<sub>2</sub>/HRP system via microplates reader.

## MATERIALS AND METHODS

### I. Reagents and Materials

All fresh vegetables were purchased from a local supermarket in Taipei, Taiwan. All Chinese herbs were obtained from local supermarket or drug store in Taipei or Taichung. ABTS (2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid)), HRP (horseradish peroxidase, type I) and L-ascorbic acid were purchased from Sigma Chemical Co. (St. Louis, MO). Hydrogen peroxide as a 35% solution was purchased from Merck (Darmstadt, Germany). All other reagents were analytical grade and were used without further purification.

### II. Instruments and Conditions

In this report, a centrifuge, Sigma 3K30, Sigma Laborzentrifugen (Harz, Germany) and a Microplate Spectrophotometer, SPECTRAMax<sup>®</sup> PLUS, Molecular Devices (Sunnyvale, CA) were used instead of a conventional spectrophotometer. The microplate reader was set up in kinetic mode with absorbance read at 414 nm every 10 sec. The microplate reader provided rapid and sensitive UV/VIS measurements of up to 96 analytes at the same time.

### III. Preparation of Sample Extracts

The extraction buffer was 6% metaphosphoric acid with 1  $\mu$ M EDTA in double deionized water. For fresh tissues, each sample was diluted with 3 equivalents of cold extraction buffer, whereas dry matter samples were diluted with a 10-times volume of extraction buffer. The samples were homogenized in a pestle and mortar, the homogenate centrifuged at 4°C, 12000 $\times$ g and the supernatant taken for antioxidant activity assay<sup>(25,26)</sup>.

### IV. Total Antioxidant Activity Assay


The ABTS/H<sub>2</sub>O<sub>2</sub>/HRP assay system for antioxidant activity was reported by Arnao *et al.*<sup>(21)</sup>. The working solution contained 2.0 mM ABTS and 0.86 nM horseradish peroxidase in 50 mM phosphate buffer, pH 7.0. The reaction was initiated with 1 mM hydrogen peroxide in 50 mM phosphate buffer.

In a 96-well microplate, each well was loaded with 240  $\mu$ L working solution, 10  $\mu$ L sample and finally 25  $\mu$ L

of H<sub>2</sub>O<sub>2</sub> to initiate the reaction. The reading was taken immediately and the time for green ABTS<sup>•+</sup> radical formation was calculated. A standard curve was plotted with different concentrations of L-ascorbic acid. Each measurement was expressed with mg L-ascorbic acid equivalent per gram fresh plant tissue.

## RESULTS AND DISCUSSION

To save time and reagents, a microplate reader was used to measure total antioxidant activity of fluid samples in this experiment. Ninety-six samples could be measured simultaneously, and it required only one-third the amount of reagents as compared to the conventional spectrophotometric method. Unlike fixed-point method, ABTS/H<sub>2</sub>O<sub>2</sub>/HRP system uses method and has absorbance readings taken every 10 sec. The plant pigments will make background of absorbance higher without influencing the lag time. This is the advantage of kinetic method.



**Figure 1.** Time course of ABTS<sup>•+</sup> radical formation in the ABTS/H<sub>2</sub>O<sub>2</sub>/HRP system using different concentrations of L-ascorbic acid as standard.


**Table 1.** Total antioxidant activities of numerous vegetables

Common name	Total antioxidant activity ( $\mu$ g/g fresh weight) <sup>a</sup>
Sugar beet root	2029 $\pm$ 77
Red cabbage leaf	1217 $\pm$ 231
Edible rape leaf	710 $\pm$ 101
Pak-choi leaf	475 $\pm$ 72
Lettuce leaf	446 $\pm$ 67
Spinach leaf	434 $\pm$ 55
Pea sprout	296 $\pm$ 44
Mungbean sprout	243 $\pm$ 56
Bitter melon fruit	228 $\pm$ 33
Carrot root	169 $\pm$ 20
Tomato fruit	143 $\pm$ 12
Head lettuce leaf	123 $\pm$ 15
Cucumber fruit	104 $\pm$ 15
Celery leaf	83 $\pm$ 4.1
Alfalfa sprout	82 $\pm$ 6.0
Soybean sprout	79 $\pm$ 5.0

<sup>a</sup>All values in this table represent the mean  $\pm$  SD (n = 3).

Figure 1 shows the characteristic time course of ABTS<sup>+</sup> radical absorption at wavelength 414 nm in ABTS/H<sub>2</sub>O<sub>2</sub>/HRP system using different concentrations of L-ascorbic acid as standard. A linear correlation was observed between lag time of green ABTS<sup>+</sup> product formation and L-ascorbic acid concentration of 50 to 1500  $\mu$ M L-ascorbic acid (Figure 2).

With the exception of red cabbage and sugar beet, total antioxidant activity of the tested vegetables ranged from 79 to 710  $\mu$ g L-ascorbic acid equivalent per gram plant tissue. Sugar beet and red cabbage had total antioxidant activity of 2029 and 1217  $\mu$ g L-ascorbic acid equivalent per gram fresh plant tissue respectively (Table 1). The high antioxidant activity of these 2 vegetables is probably due to their red pigments, flavonoids, which are very powerful antioxidants<sup>(27,28)</sup>.



**Figure 2.** Plot of ABTS<sup>+</sup> radical formation lag time versus L-ascorbic acid concentration in the ABTS/H<sub>2</sub>O<sub>2</sub>/HRP system.

**Table 2.** Total antioxidant activities of Chinese herbs and folk medicines

Common name	Latin name	Total antioxidant activity <sup>a</sup> ( $\mu$ g/g dry weight)
Tropical Almond Teriminalia Leaves	<i>Terminalia catappa</i> L.	28800 $\pm$ 9600
Moutan Radicis Cortex	<i>Paeonia suffruticosa</i> Andr. (Ranunculaceae)	17100 $\pm$ 400
Ligustri Lucidi Fructus	<i>Ligustrum lucidum</i> Ait. (Oleaceae)	6220 $\pm$ 40
Mori Folium	<i>Morus alba</i> L.(Moraceae)	5070 $\pm$ 10
Paeoniae Radix Alba	<i>Paeonia lactiflora</i> Pall. (Ranunculaceae)	4910 $\pm$ 20
Anemarrhenae Rhizoma	<i>Anemarrhena asphodeloides</i> Bge. (Liliaceae)	3990 $\pm$ 90
Rehmanniae Radix et Rhizoma	<i>Rehmannia glutinosa</i> (Gaertn.) Libosch. (Scrophulariaceae)	2420 $\pm$ 150
Ashitaba	<i>Angelica keiskei</i>	1890 $\pm$ 130
Leonuni Herba	<i>Leonurus heterophyllus</i> Sweet (Labiatae)	1630 $\pm$ 50
Asari Herba Radice	<i>Asarum heterotropoides</i> Fr.Schmidt var. <i>mandshuricum</i> (Maxim.) Kitag. (Aristolochiaceae)	1260 $\pm$ 2
Niu-chang-chih	<i>Antrodia Camphorata</i> powder	1200 $\pm$ 17
Praeparatum Mungo	<i>Phaseolus radiatus</i> L. (Leguminosae)	926 $\pm$ 62
Ligustici Rhizoma et Radix	<i>Ligusticum sinense</i> Oliv. (Umbelliferae)	901 $\pm$ 27
Mori Radicis Cortex	<i>Morus alba</i> L. (Moraceae)	850 $\pm$ 10
Angelicae Dahuricae Radix	<i>Angelica anomala</i> Lallem. (Umbelliferae)	817 $\pm$ 22
Alpiniae Oxyphyllae Fructus	<i>Alpinia oxyphylla</i> Miq. (Zingiberaceae)	568 $\pm$ 5.1
Angelicae Sinensis Radix	<i>Angelica sinensis</i> (Oliv.) Diels (Umbelliferae)	567 $\pm$ 6.0
Eucommiae Cortex	<i>Eucommia ulmoides</i> Oliv. (Eucommiaceae)	561 $\pm$ 14
Spirodela Herba	<i>Spirodela polyrrhiza</i> Schleid (Lemnaceae)	559 $\pm$ 7.0
Albiziae Cortex	<i>Albizia julibrissin</i> Durazz. (Leguminosae)	522 $\pm$ 4.1
Tribuli Fructus	<i>Tribulus terrestris</i> L. (Zygophyllaceae)	516 $\pm$ 1.0
Atractylodis Ovatae Rhizoma	<i>Atractylodes macrocephala</i> Koidz (Campositaceae)	471 $\pm$ 2.1
Cuscutae Semen	<i>Cuscuta chinensis</i> Lam. (Convolvulaceae)	466 $\pm$ 1.1
Sileris Radix	<i>Saposhnikovia divaricata</i> (Turcz.) Schischk. (Umbelliferae)	460 $\pm$ 2.1
Ophiopogonis Radix	<i>Ophiopogon japonicus</i> (L.f.) Ker-Gawl. (Liliaceae)	411 $\pm$ 7.3
Clematidis Radix	<i>Clematis chinensis</i> Osbeck (Ranunculaceae)	372 $\pm$ 3.2
Trichosanthis Frutuos	<i>Trichosanthes kirilowii</i> Maxim (Cucurbitaceae)	352 $\pm$ 8.1
Codonopsis Pilosulae Radix	<i>Codonopsis pilosula</i> (Franch.) Nannf. (Campositaceae)	291 $\pm$ 10
Platycodi Radix	<i>Platycodon grandiflorum</i> (Jacq.) A. DC.(Campanulaceae)	289 $\pm$ 1.1
Persicae Semen	<i>Prunus persica</i> (L.) Batsch (Rosaceae)	233 $\pm$ 4.0
Magnoliae Liliflorae Flos	<i>Magnolia liliflora</i> Desr. (Orchidaceae)	181 $\pm$ 13
Ampelopsitis Radix	<i>Ampelopsis japonica</i> (Thunb.) Mak. (Vitaceae)	163 $\pm$ 7.1
Taiwan Jewel Orchid	<i>Anoetochilus formosanus</i> Hayata	146 $\pm$ 31
Asparagi Radix	<i>Asparagus cochinchinensis</i> (Lour.) Merr.(Liliaceae)	145 $\pm$ 3.1
Ginseng Radix	<i>Panax ginseng</i> C.A.Mey (Araliaceae)	131 $\pm$ 5.0
Pruni Semen	<i>Prunus japonica</i> Thunb. (Rosaceae)	120 $\pm$ 14
Ginkgo Semen	<i>Ginkgo biloba</i> L. (Ginkgoaceae)	99.1 $\pm$ 11.4
Coicis Semen	<i>Coix lacryma-jobi</i> L. var.ma-yuen (Roman.) Stapf (Gramineae)	85.1 $\pm$ 7.7
Benincasae Semen	<i>Benincasa hispida</i> (Thunb.) Cogn.(Cucurbitaceae)	52.0 $\pm$ 20.6
Armeniaca Semen	<i>Prunus armeniaca</i> L. var. <i>ansu</i> Maxim (Rosaceae)	47.1 $\pm$ 8.7
Pinelliae Rhizoma	<i>Pinellia ternata</i> (Thunb.) Breit. (Araceae)	29.7 $\pm$ 6.1
Poria	<i>Poria cocos</i> (Schw.) Wolf (Polyporaceae)	21.8 $\pm$ 1.3

<sup>a</sup>All values in this table represent the mean  $\pm$  SD (n = 3).

Antioxidation activity represents the capability of scavenging free radical and offering hydrogen atom. The higher the antioxidation activity, the stronger its capability is<sup>(29)</sup>. Various methods used to evaluate total antioxidant activity of common vegetables had been reported, including oxygen radical absorption capacity method<sup>(30)</sup>, ferric reducing antioxidant capacity method<sup>(30)</sup>, liposome assay<sup>(31)</sup>, and total oxyradical scavenging capacity assay<sup>(32)</sup>. The tested samples were not overlapped with the materials used in this study, so the results cannot be compared. However, the method of total antioxidant activity determination with ABTS/H<sub>2</sub>O<sub>2</sub>/HRP system in microplate offers a fast and reliable approach.

The tissue of some Chinese herbs or folk medicines demonstrated potent antioxidative activity. For example, that of Tropical Almond Terminalia fallen leaves and Moutan Radicis Cortex were overwhelmingly high. They were 28800 and 17100  $\mu$ g L-ascorbic acid equivalent per gram weight respectively. Their high antioxidative capacity may constitute one of the beneficial factors in these traditional Chinese medication (Table 2). However, the active components that contribute to the strong antioxidant activity of these Chinese herbs is still unidentified.

Herbal drugs such as Moutan Radicis Cortex, Ligustri Lucidi Fructus, Mori Folium, Paeoniae Radix Alba, Anemarrhenae Rhizoma, Rehmanniae Radix et Rhizoma, Ashitaba, Leonuni Herba, and Asari Herba Radice were found to have higher total antioxidative activities. These drugs used to nourish qi (in Chinese, vital energy), blood, lung, and liver<sup>(33)</sup>. However, other Chinese herbs such as Asparagi Radix, Coicis Semen, Codonopsis Pilosulae Radix, and Ginseng Radix, etc. showed lower total antioxidant activities. This finding indicates that antioxidative effects are not the only beneficial factors of these medicinal herbs.

Oxidation is found to have a strong relationship with melanin formation. Antioxidants and free radical scavengers could contribute to lightening of skin pigment<sup>(34,35)</sup>. Meanwhile, skin possesses an extremely efficient antioxidant system<sup>(36)</sup> by two major groups: enzymes (e.g., superoxide dismutase, SOD) and small molecules (e.g., L-ascorbic acid). Both antioxidants could protect skin damage from oxidation. The herbs with higher antioxidant activities have the potential to be used in cosmetics.

## RRFERENCES

1. Niwa, Y. 1991. Effect of Maharishi 4 and Maharishi 5 on inflammatory mediators with special reference to their free radical scavenging effect. *Indian J. Clin. Pract.* 1: 23-27.
2. Richards, R. T. and Sharma, H. M. 1991. Free radicals in health and disease. *Indian J. Clin. Pract.* 2: 15-26.
3. Hong, C. Y., Lo, Y. C., Tan, F. C., Wei, Y. H. and Chen, C. F. 1994. *Astragalus membranaceus* and *Polygonum multiflorum* protect rat heart mitochondria against lipid peroxidation. *Am. J. Chin. Med.* 22: 63-70.
4. Hong, C. Y., Wang, C. P., Lo, Y. C. and Hsu, F. L. 1994. Effect of flavan-3-ol tannins purified from *Camellia sinensis* on lipid peroxidation of rat heart mitochondria. *Am. J. Chin. Med.* 22: 285-292.
5. Houghton, P. 1994. Ginkgo. *Pharm. J.* 253: 122-123.
6. Lo, Y. C., Teng, C. M., Chen, C. F., Chen, C. C. and Hong, C. Y. 1994. Magnolol and honokiol isolated from *Magnolia officinalis* protect rat heart mitochondria against lipid peroxidation. *Biochem. Pharmacol.* 47: 549-553.
7. Smith, P. F., MacLennan, K. and Darlington, C. L. 1996. The neuroprotective properties of the *Ginkgo biloba* leaf: a review of the possible relationship to platelet activating factor. *J. Ethnopharmacol.* 59: 131-139.
8. Burton, G. W. 1994. Vitamin E: molecular and biological function. *Proc. Nutr. Soc.* 53: 251-262.
9. Cos, P., Ying, L., Calomme, M., Hu, J. P., Cimanga, K., Poel, B. V., Pieters, L., Vlietinck, A. J. and Berghe, D. V. 1998. Structure-activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavenger. *J. Nat. Prod.* 61: 71-76.
10. Kandaswami, C. and Middleton, E. 1994. Free radical scavenging and antioxidant activity of plant flavonoids. In "Free Radicals in Diagnostic Medicine." pp. 351-376. Armstrong, D. ed. Plenum. New York, U. S. A.
11. Mantle, D., Anderton, J. G., Falkous, G., Barnes, M., Jones, P. and Perry, E. K. 1998. Comparison of methods for determination of total antioxidant status: application to analysis of medicinal plant essential oils. *Comp. Biochem. Physiol., B. Comp. Biochem.* 121: 385-391.
12. Mortensen, A. and Skibsted, L. 1997. Importance of carotenoid structure in related scavenging reactions. *J. Agric. Food Chem.* 45: 2970-2977.
13. Rouseff, R. and Nagy, S. 1994. Health and nutrition benefits of citrus fruit components. *Food Technol.* 48: 125-139.
14. Torel, J., Cillard, J. and Cillard, P. 1986. Antioxidant activity of flavonoids and reactivity with peroxy radicals. *Phytochemistry* 25: 383-385.
15. Narasimhan, R., Toshihiko, O., Hiroto, O. and Shunro, K. 1995. The contribution of plant food antioxidants to human health. *Trends Food Sci. Technol.* 6: 75-81.
16. Woodside, J. V., Young, I. S. and Yarnell, J. W. G. 1999. Fruit, vegetables and antioxidants. In "Antioxidants in Human Health and Disease." pp. 205-216. Basu, T. K., Temple, N. J. and Garg, M. L. eds. CABI Publishing. New York, U. S. A.
17. Castelluccio, C., Paganga, G., Melikian, N., Bolwell, G. P., Pridham, J., Sampson, J. and Rice-Evans, C. 1995. Antioxidant potential of intermediates in phenylpropanoid metabolism in higher plants. *FEBS Lett.* 368: 188-192.
18. Rice-Evans, C., Miller, N. J. and Paganga, G. 1997. Antioxidant properties of phenolic compounds. *Trends*



- Plant Sci. 2: 152-159.
19. Perry, L. M. 1980. Medicinal plants of east and southeast Asia-Attributed properties and uses. The Massachusetts Institute of Technology Press. p. 80.
  20. Chinese Pharmacopoeia Commission. 1999. The Coloured Atlas of Chinese Material Medica Specified in Chinese Pharmacopoeia. Guang Dong Scientific Publishing Corporation, Taiwan.
  21. Arnao, M. B., Cano, A., Hernandez-Ruiz, J., Garcia-Canovas, F. and Acosta, M. 1996. Inhibition by L-ascorbic acid and other antioxidants of the 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid) oxidation catalyzed by peroxidase: a new approach for determining total antioxidant status of foods. *Anal. Biochem.* 236: 255-261.
  22. Rice-Evans, R. and Miller, N. J. 1994. Total antioxidant status in plasma and body fluids. *Method. Enzymol.* 234: 279-293.
  23. Miller, N. J., Rice-Evans, C., Davis, M. J., Gopinathan, V. and Milner, A. 1993. A novel method for measuring antioxidant capacity and its application to monitoring the antioxidant status in premature neonates. *Chin. Sci.* 84: 407-412.
  24. Yeh, D. B. and Kuo, J. M. 2000. Simple microplate assay for determining superoxide dismutase activity. *Food Sci. Agric. Chem.* 2: 115-120.
  25. Motchnik, P. A., Frei, B. and Ames, B. N. 1994. Measurement of antioxidants in human blood plasma. *Method. Enzymol.* 234: 269-279.
  26. Behrens, W. A. and Madere, R. 1987. A highly sensitive high-performance liquid chromatography method for the estimation of ascorbic and dehydroascorbic acid in tissues, biological fluids, and foods. *Anal. Biochem.* 165: 102-107.
  27. Baublis, A., Spomer, A. and Berber-Jimenez, M. D. 1994. Anthocyanin pigments: comparison of extract stability. *J. Food Sci.* 59: 1219-1222.
  28. Vinson, J. A., Hao, Y., Su, X. and Zubik, L. 1998. Phenol antioxidant quantity and quality in foods: Vegetables. *J. Agric. Food Chem.* 46: 3630-3634.
  29. Halliwell, B. and Gutteridge, M. C. 1990. The antioxidants of human extracellular fluids. *Arch. Biochem. Biophys.* 280: 1-8.
  30. Ou, B., Huang, D., Hampsch-Woodill, M., Flanagan, J. A. and Deemer, E. K. 2002. Analysis of antioxidant activities of common vegetables employing oxygen radical absorbance capacity (ORAC) and ferric reducing antioxidant power (FRAP) assays: a comparative study. *J. Agric. Food Chem.* 50: 3122-3128.
  31. Roberts, W. G. and Gordon, M. H. 2003. Determination of the total antioxidant activity of fruits and vegetables by a liposome assay. *J. Agric. Food Chem.* 51: 1486-1493.
  32. Chu, Y. F., Sun, J., Wu, X. and Liu, R. H. 2002. Antioxidant and antiproliferative activities of common vegetables. *J. Agric. Food Chem.* 50: 6910-6916.
  33. Shanghai Science and Technology Press. 1999. The Dictionary of Chinese Medicines. Chiang-Su Medical College, Shanghai, China.
  34. Kazuhisa, M. and Minoru, F. 1991. *In vitro* effectiveness of several whitening cosmetic components in human melanocytes. *J. Soc. Cosmet. Chem.* 42: 361-368.
  35. Talwar, H. S., Griffiths, C. E., Fisher, G. J., Russman, A., Krach, K., Benrazavi, S. and Voorhees, J. J. 1993. Differential regulation of tyrosinase activity in skin of white and black individuals *in vivo* by topical retinoic acid. *J. Invest. Dermatol.* 100: 800-805.
  36. Kohen, R. 1999. Skin antioxidants: their role in aging and in oxidative stress-new approach for their evaluation. *Biomed. Pharmacother.* 53: 181-192.