

# Molecular Authentication of the Chinese Herb Huajuhong and Related Medicinal Material by DNA Sequencing and ISSR Markers

CHANG SU<sup>1</sup>, KA-LOK WONG<sup>2</sup>, PAUL PUI-HAY BUT<sup>2,4</sup>, WEI-WEI SU<sup>1</sup> AND PANG-CHUI SHAW<sup>1,2,3\*</sup>

<sup>1</sup>. School of Life Science, Sun Yat-sen University, Guangzhou, China

<sup>2</sup>. Institute of Chinese Medicine, The Chinese University of Hong Kong, Hong Kong, China

<sup>3</sup>. Department of Biochemistry, The Chinese University of Hong Kong, Hong Kong, China

<sup>4</sup>. Department of Biology, The Chinese University of Hong Kong, Hong Kong, China

(Received: November 16, 2009; Accepted: March 29, 2010)

## ABSTRACT

DNA sequences in the chloroplast (*trnH-psbA* intergenic spacer) and nuclear (ITS) regions were amplified and determined for Huajuhong derived from the peels of the immature fruits of *Citrus grandis* 'Tomentosa' and related medicinal material. These sequences together with the inter-simple sequence repeats (ISSR) markers may be used to differentiate *C. grandis* 'Tomentosa' from other *Citrus* variants for the prevention of misuse.

Key words: pomelo peel, *Citrus*, Rutaceae, authentication, *trnH-psbA*, intergenic spacer (ITS), inter-simple sequence repeats (ISSR)

## INTRODUCTION

The Chinese herb Huajuhong (*Exocarpium Citri grandis*), according to the 2005 edition of the Pharmacopoeia of the People's Republic of China, is the peel from the immature fruits of both a hairy cultivar and the common form of pomelo. The source plants, described in that Pharmacopoeia, are, respectively, *Citrus grandis* Osbeck 'Tomentosa' and *Citrus grandis* Osbeck. The correct scientific names, however, should be *Citrus maxima* (Burm.) Merr. 'Tomentosa' and *Citrus maxima* (Burm.) Merr., respectively<sup>(1-4)</sup>.

Huajuhong is a popular Chinese medicinal material for the relief of tussis and phlegm symptoms. Peel of the hairy cultivar is traditionally regarded as superior with stronger antitussive function<sup>(5,6)</sup> and thus commands a much higher market value. It differs from the common form in having densely tomentose fruit wall. The best production area of the cultivar is Huazhou city of Guangdong Province of China, so the fruit is called Huazhouyou in Chinese, while the common form is

called You (HZY and YO, respectively, hereafter).

In many herbal markets, YO which constitutes of different *Citrus* species, is often used as substitute or adulterant of HZY. The commercial pomelo peels are always in shredded slices. The dense hairy fruit wall of HZY is not a reliable character to differentiate these two commodities. On the other hand, the usefulness of chemical methods is limited since the different growth condition, storage condition, age of the sample, and processing and extracting method of the *Exocarpium Citri grandis* all affect the result<sup>(7,8)</sup>. Therefore, a reliable authentication method for *Citrus* species is essential for the prevention of misuse. Recently, DNA techniques have been developed in the area of phylogeny and authentication studies between closely related species. For examining the relationship among the *Citrus* genera, RAPD<sup>(9)</sup>, SSR<sup>(10,11)</sup>, ISSR<sup>(9,12)</sup>, SRAPs<sup>(13)</sup>, nuclear DNA<sup>(14,15)</sup> and chloroplast DNA<sup>(4,16,17)</sup> have been used. By Random Amplified Polymorphic DNA (RAPD), it has been shown that sexual reproduction and the changes of production areas can result in the genetic diversity of *Citrus grandis* "Tomentosa"<sup>(18)</sup>. It has also been reported that there are some minor differences in the ITS sequences between

\* Author for correspondence. Tel: +852-2609-6803;  
Fax: +852-2603-7246; E-mail: pcsshaw@cuhk.edu.hk

**Table 1.** Samples used in this study and GenBank accession number of the DNA sequences.

Number	Species	Sample Code	Place for collection	GenBank accession no.
1-5	<i>Citrus grandis</i> 'Tomentosa'	HZY-01 ~ HZY-05	Qinghu, Luchuan, Guangxi, China	
6-11	<i>C. grandis</i> 'Tomentosa'	HZY-06 ~ HZY-11	Daling, Pingding, Huazhou, Guangdong, China	
12	<i>C. grandis</i> 'Tomentosa'	HZY-12	Guangzhou University of Chinese Medicine, Guangdong, China	ITS: GQ231956- GQ231962 <i>trnH-psbA</i> : GQ267053-GQ267058, GQ267065
13	<i>C. grandis</i> 'Tomentosa'	HZY-13	Southern Medical University, Guangdong, China	(Note: These no. cover 1-15)
14	<i>C. grandis</i> 'Tomentosa'	HZY-14	Qinghu, Luchuan, Guangxi, China	
15	<i>C. grandis</i> 'Tomentosa'	HZY-15	Qinghu, Luchuan, Guangxi, China	
16	<i>Citrus grandis</i> Osbeck	YO-01	Mei country, Guangdong, China	
17	<i>C. grandis</i> Osbeck	YO-02	Yulin, GuangXi, China	ITS: GQ231963- GQ231965 <i>trnH-psbA</i> : GQ267059-GQ267061 (Note: These no. cover 16-19)
18	<i>C. grandis</i> Osbeck	YO-03	Guangzhou, Guangxi, China	
19	<i>C. grandis</i> Osbeck	YO-04	Jiangkou, Guiping, Guangxi, China	
20	<i>Citrus chachiensis</i>	GAN	Guangzhou University of Chinese Medicine, Guangdong, China	ITS: GQ231966 <i>trnH-psbA</i> : GQ267062
21-22	<i>Citrus reticulata</i>	JU-01-JU-02	Guangzhou University of Chinese Medicine, Guangdong, China	ITS: GQ231967 <i>trnH-psbA</i> : GQ267063
23	<i>Citrus medica</i> var. <i>sarcodactylis</i>	FO	Guangzhou University of Chinese Medicine, Guangdong, China	ITS: GQ231968 <i>trnH-psbA</i> : GQ267064

**Table 2.** Analysis of the polymorphic bands of ISSR

Primers	Sequences (5'→3')	No. of loci	No. of polymorphic loci	Polymorphism bands (%)
818	CAC ACA CAC ACA CAC AG	11	11	100
	GTG TGT GTG TGT GTG TA	8	8	100
819	ACA CAC ACA CAC ACA CG	10	8	80
827	CAC ACA CAC ACA CAC ARG	13	12	92.3
848	ACA CAC ACA CAC ACA CYT	8	8	100
855	BHB GAG AGA GAG AGA GA	7	5	71.4
885				
	Total	57	52	91.2

*Citrus grandis* "Tomentosa" and the common form, with similarity among them at 97.5%<sup>(19)</sup>.

Here, we employ chloroplast *trnH-psbA* intergenic spacer, nuclear internal transcribed spacer (ITS) and inter-simple sequence repeats (ISSR) marker to differentiate *C. grandis* 'Tomentosa' and *C. grandis*. Three other common cultivated *Citrus* species, including *C. chachiensis* (*C. reticulata* Blanco var. *chachiensis*), *C. reticulata* Blanco and *C. medica* L. var. *sarcodactylis* (Hoola van Nooten) Swingle, were also studied to prevent the misuse of these valuable herbs.

## MATERIALS AND METHODS

### I. Plant Materials

Fresh samples were collected from various sources, which were identified by Dr. W. B. Liao from Sun Yat-sen University (Guangzhou, China) according to the organoleptic characteristics (Table 1). All samples were washed with double distilled water and rinsed with 70% (v/v) ethanol to remove surface contaminants. Samples were stored in box with silica gel and kept in the School of Life Science, Sun Yat-sen University.

### II. ISSR Studies -DNA Extraction and PCR

Total DNA for ISSR study was isolated from samples using Dneasy Plant Mini Kit (Qiagen, Germany) according to the instruction of the manufacturer. ISSR amplification reactions were carried out in 25  $\mu$ L volume containing 20 ng of template DNA, 1 $\times$  *Taq* buffer [50 mM  $(\text{NH}_4)_2\text{SO}_4$ ; 75 mM Tris-HCl (pH 8.3); 50 mM KCl; 0.001% gelatin], 1 mM dNTPs, 1 unit of *Taq* polymerase and 1  $\mu$ M primers (designed by The University of British Columbia). PCR amplification was performed as follows: initial 5 minutes at 94°C, 40 cycles of 30 s at 94°C, 45 s at 55°C, 2 minutes at 72°C, and a final 7 minutes extension at 72°C. PCR amplification products were analyzed on 1.8% (w/v) agarose gel. DNA marker was prepared according to the manufacturer's instruction (Seegene, Korea).

### III. ISSR Data Analysis

The ISSR bands were scored as present (1) or absent (0), each of which was treated as an independent character regardless of its intensity. The genetic identity and genetic distance were computed using POPGENE 32, percentage of all loci that were polymorphic regardless of allele frequencies was performed by diploid data analysis of POPGENE 32. A dendrogram was constructed based on Nei's genetic distances using the un-weighted pair-group mean algorithm (UPGMA) of Molecular Evolutionary Genetic Analysis (MEGA) version 4.0<sup>(20)</sup>.

### IV. DNA Sequence Studies - DNA Extraction, PCR and Cloning

Total DNA was extracted from fresh samples according to a published method<sup>(21)</sup>. In brief, cetyltrimethyl ammonium bromide method was used for the extraction. Chloroform-isoamyl alcohol (24:1) were added to remove protein, and 2/3 (v/v) isopropanol to precipitate DNA. Finally, DNA pellet was washed with 70% ethanol, and resuspended in water. Primer ITS-5 (5'-GGAAGTA-AAAGTCGTAACAAGG-3') and ITS-4 (5'-TCCTCC-GCTTATTGATATGC-3') were used to amplify the ITS region<sup>(22)</sup>, while primer psbAF (5'-GTTATGCAT-GAACGTAATGCTC-3') and *trnHR* (5'-CGCGCATG-GTGGATTACCAAATC-3') were used to amplify *trnH-psbA* region<sup>(23)</sup>. PCR was carried out in a 25  $\mu$ L mixture containing 10 ng DNA, 1 $\times$  *Taq* buffer, 1 mM dNTPs, 1  $\mu$ M primers, and 1 unit of *Taq* polymerase. Samples were initially denatured at 94°C for 5 minutes, and then subjected to 35 PCR cycles of 94°C for 1 minute, 50°C for 1 minute and 72°C for 2 minutes. PCR products were separated on a 1.5% agarose gel. PCR products were recovered from agarose gel using the Gel-MTM Gel Extraction System (Viogene, Taiwan). Purified DNA fragment was cloned into pGEM-T Easy vector (Promega, USA). Rapid Plasmid Miniprep System (Viogene, Taiwan) was then used for plasmid purification.

### V. DNA Sequencing and Phylogenetic Analysis


Two colonies for each sample were sequenced. Primer flanking sites on the DNA sequences were removed. DNA sequences were aligned by Clustal W<sup>(24,25)</sup>. Molecular evolutionary analyses were conducted using Molecular Evolutionary Genetic Analysis (MEGA) version 4.0. Phylogenetic tree was constructed using the maximum parsimony method with default settings. Bootstrap support values were determined using 500 replicates.

## RESULTS

### I. ISSR Amplification and Phylogenetic Relationships

Totally, 76 primers were screened and six of them were capable of generating polymorphic profiles (Figure 1). ISSR primers produced varying numbers of DNA fragments, depending on their SSR motifs. Amplifications using the five 5'-anchored dinucleotide repeat ISSR primers produced an average of 5.1 bands over all the samples, among which, primers based on poly(CA) motif produced seven bands on average.

The six primers based on poly(CA) motif produced 57 bands across 23 samples, of which 52 were polymorphic bands and account for 91.23%. The number of generated bands varied from 7 (ISSR 885) to 13 (ISSR 848),




**Figure 1.** ISSR profiles of the 23 *Citrus* samples using primer (a) 818, (b) 819, (c) 827, (d) 848, (e) 855, (f) 885. M: 100 bp ladder; lane 1-15, HZY01-15; lane 16-19, YO01-04; lane 20, GAN; lane 21-22, JU and lane 23, FO.

and the size ranged from 200 to 2200 bp. The average number of bands and polymorphic bands per primer were 9.5 and 8.7, respectively. Percentage of polymorphism ranged from 71.43% (ISSR 885) to 100% (ISSR 818, 819, 855), with mean 90.62% across all samples. The 3'-anchored primer based on (GA) motifs produced a lower polymorphism rate of 71.43% (Table 2).

The ISSR bands were counted for the presence or absence among samples and the binary scores were used for the UPGMA cluster analysis. The complete data was based on a total of 57 bands. A dendrogram based on UPGMA analysis with ISSR data is shown in Figure 2. The 23 samples were grouped into two clusters. Cluster I is mainly divided into two minor clades, which consist of *C. grandis* "Tomentosa" (HZY) and *C. grandis* Osbeck (YO). These two varieties are closely related and nested in this tree, but form separate clades respectively. Cluster II consist of *C. chachiensis* (GAN), *C. reticulata* (JU) and *C. medica* var. *sarcodactylis* (FO). Contrary to *C. medica* var. *sarcodactylis* (FO), *C. chachiensis* (GAN) and *C. reticulata* (JU) are closely related.

## II. Sequence Analysis

Determined DNA sequences were deposited in GenBank with accession numbers listed in Table 1. Excluding the primer flanking site, the sizes of ITS regions (including partial 18S rRNA, ITS1, 5.8S rRNA, ITS2, and partial 26S rRNA) ranged from 701 bp to 711



**Figure 2.** Dendrogram based on the analysis of the ISSR data by UPGMA. Full names of the symbols are listed in Table 1.

HZY-01	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
HZY-08	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
HZY-11	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
HZY-12	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
HZY-13	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
HZY-14	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
HZY-15	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
YO-1	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
YO-2	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
YO-3	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
GAN	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAT	300
JU	ACGGTGCCTCGGGTGCCTCGGGCTTCTTCACTGTATCCAAACGACTCTCGGCAAC	300
FO	ACAGTGCCTACGGGTGCAGTGCCTCTTCAAATGTATCCAAACGACTCTCGGCAAC	290
***** * ***** * ***** * ***** * *****		
HZY-01	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
HZY-08	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
HZY-11	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
HZY-12	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
HZY-13	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
HZY-14	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
HZY-15	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
YO-1	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
YO-2	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
YO-3	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
GAN	GGATATCTTAGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
JU	GGATATCTCGCTCGCATCGATGAAGAACGTTGCGATACTTGGTGTGAATT	360
FO	GGATATCTCAGGCTCGTATCGATGAAGAACATAGCAAATACGATACTTGGTGTGAATT	350
***** * ***** * ***** * ***** * *****		
HZY-01	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
HZY-08	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
HZY-11	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
HZY-12	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
HZY-13	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
HZY-14	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
HZY-15	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
YO-1	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
YO-2	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
YO-3	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
GAN	AGGGCATGTCGCTTGTGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	479
JU	AGGGCAGGCTGCTGGGTGTACCGCATCGTTGCTCCACCCACCCCGCAAACCAAGGC	480
FO	AGGGCATGTCGCTGGGTGTACCGCATCGTTGCCCCACCCACCCCGCAAACCAAGGC	470
***** * ***** * ***** * ***** * *** *****		

**Figure 3.** ITS sequences of *Citrus grandis* 'Tomentosa' (HZY) and its related species. \* denotes nucleotide identical in all sequences. Nucleotides that may be used to differentiate the concerned *Citrus* species are boxed.

HZY-01	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	539
HZY-08	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	539
HZY-11	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	539
HZY-12	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	539
HZY-13	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	539
HZY-14	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	539
HZY-15	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	539
YO-1	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGGATGGCC	539
YO-2	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGGATGGCC	539
YO-3	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGGATGGCC	539
GAN	GGGGGCCCCAGGGTGGGGCATAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	539
JU	GGGGGCCCCGGGGTGGGGGGAGATTGGCCTCCGTGCGCTGACCGCTCGCGTTGGCC	540
FO	GGGGGCCCCGGGGTGGGGCATAGATTGGCCTCCGTACGCTGACTGCTCGCGTTGGCC	530
***** * *** * *** * ***** * ***** * ***** * *****		
HZY-01	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
HZY-08	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCATGCAA	658
HZY-11	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
HZY-12	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
HZY-13	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
HZY-14	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
HZY-15	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
YO-1	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
YO-2	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
YO-3	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
GAN	AGCTACCGCTGCGGCCAGTATCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	658
JU	AGCTCCCGCACGGCCGGCTCCGAGTGGGACTCTGCGACCCCTGAAGCTCCACGCAA	660
FO	AGCTCCCGCTGCGCCGGATCTCCAAGTGTGACTCTACGACCTTGAAGCTCCACGCAA	649
*** * *** * *** * *** * *** * *** * ***		

Figure 3. Continued

bp. *C. grandis* "Tomentosa", *C. grandis* Osbeck and *C. chachiensis* were both 710 bp in size. Two polymorphic sites were found between *C. grandis* "Tomentosa" and *C. grandis* Osbeck at position 534 bp and 638 bp (Figure 3). Several characteristic sites in the alignment could help to differentiate between the two *C. grandis* variants and other *Citrus* species. For example, there is a deletion in *C. medica* var. *sarcodactylis* from position 250 to 259 bp.

The sizes of *trnH-psbA* region of *C. grandis* 'Tomentosa' samples ranged between 514 and 516 bp. DNA sequences of these samples were slightly shorter than *C. reticulata* (530 bp), but longer than *C. grandis* Osbeck (512-514 bp) and *C. medica* var. *sarcodactylis* (507 bp), and similar to those of the *C. chachiensis* (516 bp). Alignment showed that this region was conserved between *C. grandis* 'Tomentosa' and *C. grandis* Osbeck

as there was only one variable site at position 355 bp. Nevertheless, we could distinguish the two *C. grandis* species from its relative species with the few insertion or deletion sites in the sequences (Figure 4).


Using the determined sequences, ITS and *trnH-psbA* phylogenetic trees were constructed. In the ITS tree, *C. grandis* (HZY and YO) and *C. chachiensis* (GAN) were closely related and resolved as a clade with bootstrap value of 100, while the rest *Citrus* species formed another clade (Figure 5). Similar clustering could be found in *trnH-psbA* tree: *C. grandis* 'Tomentosa' (HZY), *C. grandis* Osbeck (YO) and *C. chachiensis* (GAN) formed a clade with bootstrap value of 72 (Figure 6). However, sample HZY-15 was clustered with other YO samples to form a subclade, which was different from the ISSR and ITS phylogenetic trees.

HZY-01	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
HZY-08	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
HZY-11	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
HZY-12	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
HZY-13	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
HZY-14	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
HZY-15	GTCTTGCAGGCGGGTTTGAAAATAACGATCAATTCTGACCCCAGCTGGGGTC	120
YO-1	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
YO-2	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
YO-3	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
GAN	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
JU	GTCTTGTAGGCGGGTTTGAAAATAACGATCAAT-CTGACCCCAGCTGGGGTC	119
FO	GTCTTGTAGGCGGGTTTGAAAATAACGATCAATACTGACCCCAGCTGGGGTC	120
*****		
HZY-01	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	292
HZY-08	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	293
HZY-11	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	292
HZY-12	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	292
HZY-13	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	291
HZY-14	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	292
HZY-15	CGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	292
YO-1	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	292
YO-2	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	289
YO-3	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	293
GAN	TGTTGGTATGCGCTAATACTACTAATAAATTACTAAATTCTAAT-----TTTATTAT	293
JU	TGTTGGTATGCGCTAATACTACTAATAAATTAAATTCTGAATGTTATTATTTATTAT	298
FO	TGTTGGTATGCGCTAATACTACTAATAAATTAAATTCTAAT-----TTTATTAT	276
* *****		
HZY-08	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCATTTGGCTTTCA	405
HZY-11	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCATTTGGCTTTCA	404
HZY-12	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCATTTGGCTTTCA	404
HZY-13	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCATTTGGCTTTCA	403
HZY-14	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCATTTGGCTTTCA	404
HZY-15	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCAGTTGGCTTTCA	404
YO-1	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCATTTGGCTTTCA	404
YO-2	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGACCTTCATTTGGCTTTCA	401
YO-3	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCATTTGGCTTTCA	405
GAN	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCTCTCGATCTTCATTTGGCTTTCA	405
JU	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCCCTCGATCTTCATTTGGCTTTCA	418
FO	AGAAAAAAAGACAATAGAAAGGTTGAGTTTCTGCCCTCGATTTCAATTGGCTTTCA	396
*****		


**Figure 4.** *trnH-psbA* sequences of *Citrus grandis* 'Tomentosa' (HZY) and its related species. \* denotes nucleotide identical in all sequences. Nucleotides that may be used to differentiate the concerned *Citrus* species are boxed.

## DISCUSSION

ITS region of nuclear ribosomal DNA and chloroplast region *trnH-psbA* are often employed to assist identification, particularly at the intergenic level<sup>(26-28)</sup>. In this study, however, these two DNA regions could not be used to distinguish between *C. grandis* 'Tomentosa' and *C. grandis* Osbeck because there is only slight difference in the DNA sequences. Such high sequence similarity may be due to their close relationship, as also revealed by the ITS and *trnH-psbA* sequences of some species from other groups<sup>(29,30)</sup>.



**Figure 5.** Maximum parsimony tree based on the ITS sequences of *Citrus grandis* 'Tomentosa' (HZY) and its related species. *Tetradium ruticarpum* (EU663538), *C. sinensis* (AB456120), *C. aurantium* (AB456125, AB456126), *C. reticulata* (AB456092, AM398230) and *C. limon* (AB456128) were retrieved from GenBank. *T. ruticarpum* was selected as the outgroup species. Numbers above branches are bootstrap values.



**Figure 6.** Maximum parsimony tree based on *trnH-psbA* sequences of *Citrus grandis* 'Tomentosa' (HZY) and its related species. *Clausena anisata* (AM500899) and *C. aurantium* (EF590679) were retrieved from GenBank, and the former was selected as the outgroup species.

Microsatellite technique samples the whole genome, and is extremely efficient, reproducible and highly informative<sup>(10,31-33)</sup>. (AT)<sub>n</sub> is the most abundant microsatellite in plant nuclear genomes, followed by (AG)<sub>n</sub> and (AC)<sub>n</sub><sup>(10,34)</sup>. In our study, primers (CA)<sub>8</sub>G, (AC)<sub>8</sub>G, (CA)<sub>8</sub>RG, (AC)<sub>8</sub>YT, and (GT)<sub>8</sub>A generated unique fingerprints, while primer (AT)<sub>n</sub> and (AG)<sub>n</sub> produced similar fingerprints between *Citrus grandis* 'Tomentosa' and *C. grandis* Osbeck. We found that the dinucleotide repeats of the primers were easier to produce fingerprint than that of trinucleotide repeats in *Citrus* genomic DNA, probably due to the abundance of the dinucleotide repeats in the *Citrus* genome. This is also consistent with the surveys of microsatellite markers in the sweet orange (*Citrus sinensis* L. Osbeck)<sup>(10)</sup>. We also found that only BHB(GA)<sub>7</sub> generated excellent results in ISSR, while other primers with different number of GA-repeat produced no PCR products. Poor result may be due to either the characteristics of the primers or to the relative abundance of the priming sites in the genome<sup>(35)</sup>.

HZY-15 has less hairs than the other *Citrus grandis* 'Tomentosa' samples, and people usually call this type of *C. grandis* Fu Mao Huazhouyou, which means less tomentum on the fruit surface<sup>(36)</sup>. Instead of clustering with other HZY samples, it forms a subclade with other YO samples in the *trnH-psbA* tree. On the other hand, Fu Mao-HZY and HZY cannot be distinguished by the ITS region identification as well as the ISSR UPGMA tree. This indicates that the genetic difference between Fu Mao-HZY and HZY is small.

The most widely accepted taxonomic system for *Citrus* was proposed by Swingle<sup>(37)</sup> and Tanaka<sup>(38)</sup>. Subsequent phylogenetic analysis by Barrett and Rhodes suggested that there were only three true species within cultivated *Citrus*, including citron (*Citrus medica* L.), mandarin (*C. reticulata* Blanco) and pomelo (*C. grandis*)<sup>(39)</sup>. Some cultivated *Citrus* species are used as medicinal materials for centuries in China. According to traditional Chinese medicine, all these *Citrus* variants have therapeutic effects in reducing phlegm and smoothing coughs, but are used to treat different syndromes. The *Citrus* crude drugs could be misused easily when just considering their morphological and chemical characteristics, and thus compromising the pharmaceutical efficacy. In this work, we have shown the DNA sequences in *trnH-psbA* and ITS regions for the determination of pomelo and related medicinal material. ISSR fingerprint analysis is able to differentiate between *Citrus grandis* 'Tomentosa' from other *Citrus* variants to prevent the misuse of the *Citrus* herbs. Our study also shows that ISSR and DNA sequencing methods are complementary for the differentiation of closely related species in general.

## ACKNOWLEDGMENT

This work was supported in part by a Focused

Investment Scheme Funding (Scheme A) of the Chinese University of Hong Kong.

## REFERENCES

1. Scora, R. W. and Nicolson, D. H. 1986. The correct name for the shaddock, *Citrus maxima*, not *C. grandis* (Rutaceae). *Taxon* 35: 592-595.
2. Mabberley, D. J. 2004. *Citrus* (Rutaceae): a review of recent advances in etymology, systematics and medical applications. *Blumea* 49: 481-498.
3. Mabberley, D. J. 1997. A classification for edible *Citrus* (Rutaceae). *Telopea (Syd.)* 7: 167-172.
4. Bayer, R. J., Mabberley, D. J., Morton, C., Miller, C. H., Sharma, I. K., Pfeil, B. E., Rich, S., Hitchcock, R. and Sykes, S. 2009. A molecular phylogeny of the orange subfamily (Rutaceae: Aurantioideae) using nine cpDNA sequences. *Am. J. Bot.* 96: 668-685.
5. Li, P. B., Ma, Y., Wang, Y. G. and Su, W. W. 2006. Experimental studies on antitussive, expectorant and antiasthmatic effects from *Citrus grandis* var. *tomentosa*. *Zhongguo Zhong Yao Za Zhi* 16: 1350-1352.
6. Zhang, X. M., Chen, Z. X. and Lin, L. 2004. Comparison on dissolving sputum and anti-inflammation of "Mao Ju Hong" and "Guang Ju Hong". *Zhong Yao Cai* 27: 122-123.
7. Lin, L., Huang, L. Z., Ou, J. F. and Chen, K. 2006. Observation on dynamic changes of flavonoids contents of *Citrus grandis* (L.) Osbeck var. *tomentosa* Hort. *J. Guangzhou Uni. Trad. Chin. Med.* 23: 256-261.
8. Huang, L. Z., Liang, Z. H., Lin, Li., Gao, D. X. and Ou, J. F. 2005. Influence of different processing technique on the content of naringin in Exocarpium Citrus Grandis. *Trad. Chin. Drug Res. Clin. Pharmacol.* 16: 59-61.
9. Pasquale, D. F., Siragusa, M., Abbate, L., Tusa, N., Pasquale, D. C. and Alonzo, G. 2006. Characterization of five sour orange clones through molecular markers and leaf essential oils analysis. *Sci. Hortic.* 109: 54-59.
10. Novelli, V. M., Cristofani, M., Souza, A. A. and Machado, M. A. 2006. Development and characterization of polymorphic microsatellite markers for the sweet orange (*Citrus sinensis* L. Osbeck). *Genet. Mol. Biol.* 29: 90-96.
11. Pang, X. M., Hu, C. G. and Deng, X. X. 2003. Phylogenetic relationships among *Citrus* and its relatives as revealed by SSR markers. *Chin. J. Genet.* 30: 81-87.
12. Shahsavar, A. R., Izadpanah, K., Tafazoli, E. and Tabatabbaei, S. B. E. 2007. Characterization of *Citrus* germplasm including unknown variants by inter-simple sequence repeat (ISSR) markers. *Sci. Hortic.* 112: 310-314.
13. Uzun, A., Yesiloglu, T., Aka-Kacar, Y., Tuzcu, O. and Gulsen, O. 2009. Genetic diversity and relationship within *Citrus* and related genera based on sequence related amplified polymorphism markers (SRAPs). *Sci. Hortic.* 121: 306-312.
14. Nicolosi, E., Deng, Z. N., Gentile, A., LaMalfa, S., Continella, G. and Tribulato, E. 2000. *Citrus* phylogeny and genetic origin of important species as investigated by molecular markers. *Theor. Appl. Genet.* 100: 1155-1166.
15. Deng, Z., Malfa, S. L., Xie, Y. M., Xiong, X. Y. and Gentile, A. 2007. Identification and evaluation of chloroplast uni- and trinucleotide sequence repeats in *Citrus*. *Sci. Hortic.* 111: 186-192.
16. de Araújo, E.F., de Queiroz, L.P. and Machdo M. AM. 2003. What is *Citrus*? Taxonomic implications from a study of cp-DNA evolution in the tribe Citreae (Rutaceae subfamily Aurantioideae). *Org. Divers. Evol.* 3: 55-62.
17. Jung, Y. H., Kwon, H. M., Kang, S. H., Kang, J. H. and Kim, S. C. 2005. Investigation of the phylogenetic relationships within the genus *Citrus* (Rutaceae) and related species in Korea using plastid *trnL-trnF* sequences. *Sci. Hortic.* 104: 179-188.
18. Lin, L., Ou, J. F., Xiao, F. X. and Liu, X. H. 2008. Random amplified polymorphic DNA analysis of germplasm resources of Exocarpium Citri Grandis. *J. Guangzhou Uni. Trad. Chin. Med.* 25: 350-354.
19. Chen, X. Y., Gao, X. X., Luo, Y. S., Cai, Y. W. and Wu, Y. Y. 2007. Preliminary study on rDNA ITS sequencing and characteristics of *Citrus grandis* var. *tomentosa*. *Zhong Yao Cai* 30: 268-270.
20. Tamura, K., Dudley, J., Nei, M. and Kumar, S. 2007. MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* 24: 1596-1599.
21. Kang, H. W., Cho, Y. G., Yoon, U. H. and Eun, M. Y. 1998. A rapid DNA extraction method for RFLP and PCR analysis from a single dry seed. *Plant Mol. Biol. Rep.* 16: 90.
22. Innis, M. A., Gelfand, D. H., Sninsky, J. J. and White, T. J. 1990. PCR protocols: a guide to methods and applications. Academic Press. San Diego, U. S. A.
23. Sang, T., Crawford, D. J. and Stuessy, T. 1997. Chloroplast DNA phylogeny, reticulate evolution, and biogeography of *Paeonia* (Paeoniaceae). *Am. J. Bot.* 84: 1120-124.
24. Thompson, J. D., Higgins, D. G. and Gibson, T. J. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22: 4673-4680.
25. Hall, T. A. 1999. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* 41: 95-98.
26. Baldwin, B. G., Sanderson, M. J., Porter, J. M., Wojciechowski, M. F., Campbell, C. S. and Donoghue, M. J. 1995. The ITS region of nuclear ribosomal DNA:

a valuable source of evidence on angiosperm phylogeny. *Ann. Mo. Bot. Gard.* 82: 247-277.

27. Baldwin, B. G. 1993. Molecular phylogenetics of *Calycadenia* (Compositae) based on ITS sequences of nuclear ribosomal DNA: Chromosomal and morphological evolution reexamined. *Am. J. Bot.* 80: 222-238.

28. Baldwin, B. G. 1992. Phylogenetic utility of the internal transcribed spacers of nuclear ribosomal DNA in plants: An example from the Compositae. *Mol. Phylogen. Evol.* 1: 3-16.

29. Kyndt, T., Van Droogenbroeck, B., Romeijn-Peeters, E., Romero-Motochi, J. P., Scheldeman, X., Goetghebeur, P., Van Damme, P. and Gheysen, G. 2005. Molecular phylogeny and evolution of Caricaceae based on rDNA internal transcribed spacers and chloroplast sequence data. *Mol. Phylogen. Evol.* 37: 442-459.

30. Zhao, H. G., Zhou, J. J., Cao, S. S., Zheng, Y. H., Shan, Y. and Xia, B. 2009. Analysis of interspecific relationship among *Stellaria media* and its related species based on ITS and *trnL*-F sequence differences. *J. Plant Resour. Environ.* 18: 1-5.

31. Zane, L., Bargelloni, L. and Patarnello, T. 2002. Strategies for microsatellite isolation: a review. *Mol. Ecol.* 11: 1-16.

32. Viruel, M. A. and Hormaza, J. I. 2004. Development, characterization and variability analysis of microsatellites in lychee (*Litchi chinensis* Sonn., Sapindaceae). *Theor. Appl. Genet.* 108: 896-902.

33. He, C., Poysa, V. and Yu, K. 2003. Development and characterization of simple sequence repeat (SSR) markers and their use in determining relationships among *Lycopersicon esculentum* cultivars. *Theor. Appl. Genet.* 106: 363-373.

34. Wang, Z., Weber, J. L., Zhong, G. and Tanksley, S. D. 1994. Survey of plant short tandem DNA repeats. *Theor. Appl. Genet.* 88: 1-6.

35. Fang, D. Q. and Roose, M. L. 1997. Identification of closely related *Citrus* cultivars with inter-simple sequence repeat markers. *Theor. Appl. Genet.* 95: 408-417.

36. Wu, W. Z., Chen, H. P., Ma, X. J. and Zeng, R. S. 2002. GC-MS analysis of *Citrus grandis* (L.) Osbeck var. *tomentosa*. *Zhong yao cai* 25: 180-181.

37. Swingle, W. T. and Reece, P. C. 1967. The Botany of *Citrus* and its wild relatives. In "The *Citrus* industry". pp. 190-430. University of California. Berkeley, U. S. A.

38. Tanaka, T. 1977. Fundamental discussion of *Citrus* classification. *Stud. Citrol.* 14: 1-6.

39. Barrett, H. C. and Rhodes, A. M. 1976. A numerical taxonomic study of affinity relationships in cultivated *Citrus* and its close relatives. *Syst. Bot.* 1: 105-136.