

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.jfda-online.com](http://www.jfda-online.com)

## Original Article

# Bioactive polyphenols in kinkéliba tea (*Combretum micranthum*) and their glucose-lowering activities



Cara Welch <sup>a,b,1,2</sup>, Jing Zhen <sup>a,b,1</sup>, Emmanuel Bassène <sup>c</sup>, Ilya Raskin <sup>a</sup>,  
James Edward Simon <sup>a,b,\*</sup>, Qingli Wu <sup>a,b,\*\*</sup>

<sup>a</sup> The New Use Agriculture and Natural Plant Products Program, Department of Plant Biology, School of Environmental and Biological Sciences, Rutgers University, 59 Dudley Road, New Brunswick, NJ 08901, USA

<sup>b</sup> Department of Medicinal Chemistry, Ernest Mario School of Pharmacy, Rutgers University, 160 Frelinghuysen Road, Piscataway, NJ 08854, USA

<sup>c</sup> Department of Pharmacognosy and Botany, Faculty of Medicine, Pharmacy and Odontology, UCAD, BP 5281 Dakar Fann, Senegal

## ARTICLE INFO

## Article history:

Received 26 January 2017

Received in revised form

19 April 2017

Accepted 27 May 2017

Available online 13 June 2017

## Keywords:

Kinkéliba

*Combretum micranthum*

Flavonoid

Anti-diabetes

## ABSTRACT

Herbal tea kinkéliba prepared from the leaves of *Combretum micranthum* has been widely consumed in West African countries for its flavor, nutritional and medicinal properties. Under bio-guided screening, the kinkéliba leaves were chemically investigated using various chromatographic and spectrometric methods that led to the identification of thirteen different flavonoid compounds. Further biological tests illustrated that the identified compounds may have synergistic effects to decrease the expression of phosphoenolpyruvate carboxykinase (PEPCK) mRNA and glucose production in an H4IIE hepatoma cell line, indicating its potential use for insulin-resistant diabetes treatment. Further *in vivo* study on C57BL/6J mice indicates that kinkéliba can lower plasma glucose levels in a dose-dependent manner without significant weight loss and toxicity. The ethyl acetate extract in rich of flavonoids could also increase the glucose tolerance (GT) after seven weeks' administrations. Both *in vitro* and *in vivo* experiments support a potential new application of kinkéliba leaves as an anti-diabetes agent.

Copyright © 2017, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

*Combretum micranthum* (kinkéliba) is an undomesticated shrub species found in the Tiger bush region of western Africa. It is a

bushy shrub or creeper that can reach up to 20 m in height, and common on cultivated and fallow ground, dominant in sub-Saharan Africa, with higher concentrations in Senegal, Mali, and Burkina Faso [1]. The leaves are harvested from wild growing populations and used as popular traditional herbal

\* Corresponding author. Department of Plant Biology, Rutgers University, 59 Dudley Road, New Brunswick, NJ 08901, USA.

\*\* Corresponding author. Department of Plant Biology, Rutgers University, 59 Dudley Road, New Brunswick, NJ 08901, USA.

E-mail addresses: [jimsimon@rutgers.edu](mailto:jimsimon@rutgers.edu) (J.E. Simon), [qlwu@sebs.rutgers.edu](mailto:qlwu@sebs.rutgers.edu) (Q. Wu).

<sup>1</sup> These two authors contribute equally to this work.

<sup>2</sup> Present Addresses: Division of Dietary Supplement Programs, CFSAN/FDA, 5100 Paint Branch Parkway, College Park, MD 20740, USA. <http://dx.doi.org/10.1016/j.jfda.2017.05.009>

1021-9498/Copyright © 2017, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

tea in several tropical West African savannah countries. Kinkéliba consumed as bush tea, has a pleasant flavor, light to dark green-brown color [2]. The tea is also used in traditional medicine as a general panacea and for diuretic and digestion purposes including gastrointestinal problems, colic, and vomiting [1,3–5]. Previous phytochemical studies indicated that the leaf extracts contain flavonoids including vitexin, isovitexin, orientin, homoorientin, myricetin-3-O-glucoside, and myricetin-3-O-rutinoside; alkaloids including stachydrine, hydroxyl-stachydrine, and choline; and sugar alcohols including *m*-inositol and sorbitol; and flavan alkaloids including kinkélolds A, B, C and D [5–8].

In this investigation, we originally set out to isolate and identify the polyphenols in kinkéliba that we hypothesized may be responsible for its traditional activity through solvent participation and various chromatographic and spectrometric methods including normal pressure chromatography, HPLC/UV/MS and nuclear-magnetic resonance (NMR). Yet, as we also had available a robust series of assays to screen for anti-diabetes activity, we subjected the separated compounds and extracts/fractions to a variety of these assays. The *in vitro* assays chosen for the bioactivity test of kinkéliba regarding glucose-lowering activity are measuring the percent decrease of glucose production and the down-regulation of PEPCK mRNA in rat hepatoma cells (H4IIE) [9]. A diet-induced obesity (DIO) C57BL/6J mouse model was used to further understand the anti-hyperglycemic effect of kinkéliba *in vivo*. It is a particularly good example of the human metabolic disorder because it develops a syndrome of obesity, accompanied by hyperinsulinemia and hyperglycemia (indicators of Type 2 diabetes), when allowed unfettered access to a high-fat diet [10]. The development of insulin resistance, hyperglycemia, and obesity parallel the progression of Type II diabetes in humans. We hypothesized that kinkéliba's purported health promoting properties may be in part due to the presence of polyphenols, and whereas some polyphenols have been shown in other plant species to exhibit anti-diabetic activity, we postulated that leaves of this plant may then also show some positive activity in the treatment of diabetes.

Diabetes mellitus is a chronic disease that affects 29.1 million people or 9.3% of total population in US in 2014 [11]. Specifically, Type 2 diabetes, compromising approximately 90%–95% of all diagnosed cases, often develops from excess body weight and physical inactivity resulting in the body's ineffective use of insulin, which creates hyperglycemia. There are many factors, including environmental and genetic components, contribute to the hyperglycemia of diabetes [12]. Even if the reasons are not fully understood yet, a prominent defect is that the insulin could not efficiently inhibit hepatic glucose production (HGP) [13]. Increased HGP may be due to increased gluconeogenesis, which involves a series of biochemical reactions catalyzed by several enzymes under the tight regulation of insulin [14]. The first, and also the rate-limiting step of hepatic gluconeogenesis, is controlled by enzyme phosphoenolpyruvate carboxykinase (PEPCK), which converts compound oxaloacetate to phosphoenolpyruvate. Excreted insulin is supposed to strongly repress the expression of PEPCK mRNA through the activation of enzyme phosphoinositide-3 kinase (PI3K) pathway, resulting decreased *de novo* glucose synthesis [15]. However, in diabetes

patients, insulin does not effectively inhibit the expression of PEPCK, therefore, resulting in increased blood glucose or hyperglycemia. Phytochemicals or plant extract that can suppress the expression of PEPCK mRNA, may overcome insulin resistance, and lower the glucose production from gluconeogenesis [9]. Therefore, the focus of this research shifted to determine whether bioactive polyphenols in kinkéliba exhibit anti-diabetic activity *in vitro* and *in vivo*.

## 2. Materials and methods

### 2.1. Chemicals and biochemicals

8-(4-Chlorophenylthio)-cAMP, dexamethasone, sodium pyruvate, sodium lactate, and ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid)) were purchased from Sigma–Aldrich (St. Louis, MO). Human insulin was purchased from Eli Lilly (Indianapolis, IN). Reagents used in RT-PCR, including the enzymes, were purchased from Stratagene (La Jolla, CA). The H4IIE cell line (ATCC CRL-1548) was provided by the American Type Culture Collection (Manassas, VA).

### 2.2. Plant material

Kinkéliba samples utilized for this study was collected from shrubs within a localized population in Leen, Pout, in central Senegal. Plant materials were gathered by the NGO ASNAPP-Senegal with members of the community and with prior written consent from the community and community leaders. Locally, communities often dry the leaves and branch together forming a long bundle and then sold locally as an herbal tea. The leaves and branches were manually harvested, air-dried under shade in ambient conditions and off the ground on white tarp. Once dried, the leaves were manually separated from the branches, and cleaned. The trees from which the leaves were harvested were taxonomically identified as *C. micranthum* by the authors who have been working with this same plant population and species in Senegal with the community as an indigenous herbal tea to provide additional local income opportunities. The dried leaf material were originally air-shipped to South Africa where the leaves were processed by Cape Natural Teas, Cape Town, South Africa and packaged into a Mpuntu line of African herbal teas and then air-shipped to Rutgers University for analysis. Additional dried leaves from same plant populations were air-shipped directly from Dakar, Senegal to Rutgers University for analysis.

### 2.3. General methods

1D and 2D NMR spectroscopy was performed on a Bruker Avance 400 MHz spectrometer (Billerica, MA). All organic solvents used for chromatographic separation are HPLC grade (Fisher Scientific, Springfield, NJ). Column chromatography was performed using silica gel (230–400 mesh; Selecto Scientific, Suwanee, GA), Sephadex LH-20 (25–100  $\mu$ m, Sigma-Aldrich, St. Louis, MO) and Polyamide CC 6 (50–160  $\mu$ m, Sorbent Technologies, Atlanta, GA). HPLC grade water was prepared using a Millipore Milli-Q purification system (Millipore Corp, Bedford, MA) and used for all experimental procedures.

Preparative HPLC was performed on a Waters 600/2487 system with a UV detector and Microsorb C<sub>18</sub> column (Varian, 10  $\mu$ m, 41.4  $\times$  300 mm). Analytical HPLC was performed on Waters 2695/2996 (Milford, MA) with diode array detector (DAD). LC-MS detection was performed on an Agilent 1100 series (Waldbronn, Germany) with DAD detector, MSD trap with an electrospray ion (ESI) source under positive mode. Analytical column is Microsorb C<sub>18</sub> column (Varian, 5  $\mu$ m, 4.6  $\times$  250 mm).

## 2.4. Extraction and isolation

The dried leaves (1.0 kg) were ground to fine powder and extracted with ethanol two times and a third time with an 80% ethanol (v/v) solution by maceration for 24 h at room temperature. The filtrates were combined and concentrated under reduced pressure. The crude extract was dissolved in water/ethanol (95:5, v/v) and partitioned between hexane (labeled as LH), chloroform (LC), ethyl acetate (LE), and *n*-butanol (LB), with the remaining water fraction as LW. Antioxidant activity of each extract was measured through ABTS radical scavenge assay and ethyl acetate fraction (LE) showed superior activity than other fractions (Supplemental Material S1), and followed by *n*-butanol fraction (LB), on which further investigation was conducted and with those results patented [8]. Antioxidant activities of plant extracts are often from the functional phytochemical polyphenols, which have been reported with an extensive range of biological activities [16–18]. Some of these polyphenol compounds have also been reported to exhibit *in vitro* and *in vivo* anti-diabetic activities [17,19–21]. Therefore, based on these relationships, we hypothesized that the fractions with potent antioxidant activity may also present potential glucose-lowering activity. Ethyl acetate (LE) extract was evaporated *in vacuo* to give residue (14.40 g), which was subjected to silica gel chromatographic column (5%–100% methanol/dichloromethane with 0.1% acetic acid) to give 12 fractions; labeled as LE-1 to LE-12. The fifth fraction, LE-5 (3.67 g), was further separated by Sephadex LH-20 column using 100% methanol into 9 fractions labeled as LE-5-1 to LE-5-9. LE-5-6 (700.9 mg) fraction was further purified by prep-HPLC (0%–100% acetonitrile/water with 0.1% formic acid) on Microsorb C<sub>18</sub> column to give compounds 1–4. LE-10 (4.46 g) was applied to polyamide column and eluted with a gradient of methanol/water (20–100%) to give 15 fractions, labeled as LE-10-1 to LE-10-15. Compound 5–8 were precipitated out from fraction LE-10-1, LE-10-2, LE-10-3 and LE-10-4, respectively. The concentrated LE-10-6 (166.7 mg) was then applied on prep-HPLC (0%–100% acetonitrile/water with 0.1% formic acid) with column Microsorb C<sub>18</sub> to give compound 9. The mixture of compounds 10–13 were further separated using analytical HPLC on Polaris Amide C<sub>18</sub> column (0%–100% acetonitrile/water with 0.1% formic acid) from LE-10-8 and LE-10-8.

## 2.5. Structure elucidation of compounds 1–13

The structures of compounds 1–13 were elucidated using different spectrometric methods including UV, MS and NMR spectroscopy and the spectral data was compared with earlier reports and some were further confirmed by comparison with authentic standards on HPLC. Their spectral data is described below and detailed identification is described in Section 3.1.

(-)-Epigallocatechin (1): UV  $\lambda_{\max}$ : 270 nm; ESI-MS: *m/z* 307 [M+H]<sup>+</sup>, 139 (RDA fragment, <sup>1,3</sup>A<sup>+</sup>); <sup>1</sup>H NMR (400 M Hz, MeOD<sub>4</sub>) and <sup>13</sup>C NMR (100 M Hz, MeOD<sub>4</sub>): Supplementary Material S6;

(-)-Epicatechin (2): UV  $\lambda_{\max}$ : 272 nm. ESI-MS: *m/z* 291 [M+H]<sup>+</sup>, 139 (RDA fragment, <sup>1,3</sup>A<sup>+</sup>); <sup>1</sup>H NMR (400 M Hz, MeOD<sub>4</sub>) and <sup>13</sup>C NMR (100 M Hz, MeOD<sub>4</sub>): Supplementary Material S6;

3', 4', 5', 5, 7- pentahydroxyflavan (3): UV  $\lambda_{\max}$ : 275 nm. ESI-MS: *m/z* 291 [M+H]<sup>+</sup>, 139 (RDA fragment, <sup>1,3</sup>A<sup>+</sup>); <sup>1</sup>H NMR (400 M Hz, MeOD<sub>4</sub>) and <sup>13</sup>C NMR (100 M Hz, MeOD<sub>4</sub>): Supplementary Material S6;

3', 4', 5, 7- tetrahydroxyflavan (4): UV  $\lambda_{\max}$ : 280 nm. ESI-MS: *m/z* 275, [M+H]<sup>+</sup>, 139 (RDA fragment, <sup>1,3</sup>A<sup>+</sup>); <sup>1</sup>H NMR (400 M Hz, MeOD<sub>4</sub>) and <sup>13</sup>C NMR (100 M Hz, MeOD<sub>4</sub>): Supplementary Material S6;

Vitexin (5): UV  $\lambda_{\max}$ : 216, 269, 338 nm. ESI-MS: *m/z* 433 [M+H]<sup>+</sup>, 283, 313, 397;

Isovitexin (6): UV  $\lambda_{\max}$ : 216, 269, 338 nm. ESI-MS: *m/z* 433 [M+H]<sup>+</sup>, 283, 313, 397;

Orientin (7): UV  $\lambda_{\max}$ : 216, 269, 338 nm. ESI-MS: *m/z* 449 [M+H]<sup>+</sup>, 299, 329, 359;

Homoorientin (8): UV  $\lambda_{\max}$ : 216, 269, 338 nm. ESI-MS: *m/z* 449 [M+H]<sup>+</sup>, 299, 329, 359;

Myricetin-3-O-glucoside (9): <sup>1</sup>H NMR (400 M Hz, MeOD<sub>4</sub>) characteristic peaks:  $\delta$  6.18 (1H, d, *J* = 2.0 Hz, H-6), 6.36 (1H, d, *J* = 2.0 Hz, H-8), 7.29 (2H, s, H-1', 6'), 5.23 (1H, d, *J* = 7.7 Hz, H-1''), 3.84 (1H, dd, *J* = 7.7, 8.3 Hz, H-2''), 3.51 (1H, t, *J* = 8.3 Hz, H-3''), 3.41 (1H, t, *J* = 8.3 Hz, H-4''), 3.22 (1H, m, H-5''), 3.76, 3.84 (2H, dd, H-6''); <sup>13</sup>C NMR (100 M Hz, MeOD<sub>4</sub>) characteristic peaks:  $\delta$  99 (C-6), 94 (C-8), 108 (C-2', 6'), 103 (C-1''), 76 (C-3''), 69 (C-4''), 77 (C-5''), 62 (C-6'').

2''-O-galloylvitexin (10): UV  $\lambda_{\max}$ : 216, 269, 338 nm. ESI-MS: *m/z* 585 [M+H]<sup>+</sup>, 313, 415, 433, 465. <sup>1</sup>H NMR (400 M Hz, MeOD<sub>4</sub>)  $\delta$  6.41–6.60 (1H, s, H-3), 6.07–6.22 (1H, s, H-6), 7.81, 8.06 (2H, d, *J* = 8.7 Hz, H-2', 6'), 6.96 (2H, d, *J* = 8.7 Hz, H-3', 5'), 5.20, 5.32 (1H, d, *J* = 10 Hz, H-1''), 5.62, 5.71 (1H, t, *J* = 10 Hz, H-2''), 3.84 (1H, t, *J* = 9.2 Hz, H-3''), 3.76 (1H, t, *J* = 9.2 Hz, H-4''), 3.56 (1H, m, H-5''), 3.85, 4.02 (2H, H-6''), 6.89, 6.69 (2H, s, H-2'''). <sup>13</sup>C NMR (100 M Hz, MeOD<sub>4</sub>)  $\delta$  166.75 (C-2), 103.76 (C-3), 184.20 (C-4), 162.79 (C-5), 99.19 (C-6), 164.23 (C-7), 103.76 (C-8), 158.57 (C-9), 105.69 (C-10), 123.83 (C-1'), 130.19 (C-2'), 117.05 (C-3'), 162.79 (C-4'), 117.05 (C-5'), 130.19 (C-6'), 72.56 (C-1''), 74.19 (C-2''), 78.09 (C-3''), 73.38 (C-4''), 83.25 (C-5''), 63.14 (C-6''), 121.42 (C-1'''), 110.34 (C-2'''), 146.27 (C-3'''), 139.70 (C-4'''), 146.27 (C-5'''), 110.34 (C-6'''), 167.65 (C-7''')

2''-O-galloylisorientin (11): UV  $\lambda_{\max}$ : 216, 269, 338 nm. ESI-MS: *m/z* 585 [M+H]<sup>+</sup>, 313, 415, 433, 465; <sup>1</sup>H NMR (400 M Hz, MeOD<sub>4</sub>)  $\delta$  6.48 (1H, s, H-3), 6.39 (1H, s, H-8), 7.74 (2H, d, *J* = 8.7 Hz, H-2' 6'), 6.87 (2H, d, *J* = 8.7 Hz, H-3', 5'), 5.16 (1H, d, *J* = 10 Hz, H-1''), 5.77 (1H, H-2''), 3.76 (1H, t, *J* = 9.3 Hz, H-3''), 3.62 (1H, t, *J* = 9.3 Hz, H-4''), 3.51 (1H, m, H-5''), 3.80, 3.94 (2H, H-6''), 6.94 (2H, s, H-2''', 6'''); <sup>13</sup>C NMR (100 M Hz, MeOD<sub>4</sub>)  $\delta$  166.18 (C-2), 103.92 (C-3), 183.87 (C-4), 162.70 (C-5), 108.02 (C-6), 164.50 (C-7), 158.81 (C-8), 104.97 (C-10), 123.18 (C-1'), 129.46 (C-2'), 116.99 (C-3'), 162.70 (C-4'), 116.99 (C-5'), 129.46 (C-6'), 73.87 (C-1''), 73.98 (C-2''), 78.18 (C-3''), 71.87 (C-4''), 82.88 (C-5''), 62.86 (C-6''), 121.51 (C-1'''), 110.34 (C-2'''), 146.23 (C-3'''), 139.64 (C-4'''), 146.23 (C-5'''), 110.34 (C-6'''), 167.61 (C-7''')

2''-O-galloylorientin (12): UV  $\lambda_{\max}$ : 216, 269, 338 nm. ESI-MS: *m/z* 601 [M+H]<sup>+</sup>, 329, 431, 449, 481

2''-O-galloylhomoorientin (**13**): UV  $\lambda_{\text{max}}$ : 216, 269, 338 nm. ESI-MS:  $m/z$  601  $[M + H]^+$ , 329, 431, 449, 481

## 2.6. Cell culture

The H4IIE hepatoma cells (ATCC CRL-1600) were routinely cultured in Dulbecco's modified Eagle's medium supplemented with 2.5% (v/v) newborn calf serum, 2.5% (v/v) fetal calf serum and grown at 37 °C in humidified 5% CO<sub>2</sub> atmosphere.

## 2.7. RNA extraction and quantitative real-time polymerase chain reaction (qPCR)

The H4IIE cells were plated in 24-well tissue culture plates (Greiner Bio One, Monroe, NC) and were grown to near confluence. A treatment of 500 nM dexamethasone and 0.1 mM 8-CTP-cAMP (Dex-cAMP) for 8 h was used to induce PEPCK gene expression. The cells were co-treated with plant extract (100 µg/mL), subfractions (100 µg/mL), isolated compounds (20 µM) or insulin (10 nM).

Total RNA was extracted from treated H4IIE cells using Trizol reagent (Invitrogen), treated with DNase I (Invitrogen) to remove any traces of DNA contamination, and quantitated spectrophotometrically using the NanoDrop system (NanoDrop Technologies). Gel electrophoresis was used to assess the quality of RNA sample. 2.5 µg RNA from each sample was used as template to synthesize cDNA using Stratascript reverse transcriptase (Stratagene, La Jolla, CA), following the manufacturer's protocol [9].

PCR reaction was comprised of 5 µL of each 4-fold diluted cDNA sample, 0.5 µL of 6 µM primers (IDT, Coralville, IA), and 12.5 µL of Brilliant SYBR Green PCR master mix (2X; Stratagene) supplemented with green jump-start Taq polymerase. ROX (Stratagene) was used as a reference dye. qPCR were performed on MX3000p system (Stratagene) with the following steps: initiation (1 cycle at 50 °C for 2 min, 1 cycle of 95 °C for 10 min), amplification (40 cycles of 15 s at 95 °C and 1 min at 60 °C), and dissociation (1 cycle at 95 °C for 1 min, 55 °C for 30 s, and 95 °C for 30 s). Non-template control and non-RT control were performed in each experiment to rule out the possible contamination. The primers were designed using the Primer Express software (Applied Systems, v. 2.0, Foster City, CA) as follows:  $\beta$ -actin: 5'-GGG AAA TCG TGC GTG ACA TT-3' (sense) and 5'-GCG GCA GTG GCC ATC TC-3' (anti-sense); PEPCK: 5'-GCA GAG CAT AAG GGC AAG GT-3' (sense) and 5'-TTG CCG AAG TTG TAG CCA AA-3' (anti-sense). The relative mRNA expression of PEPCK were analyzed using the DDC<sub>T</sub> method with  $\beta$ -actin as a house-keeping gene as previously described [9]. The PEPCK gene expression of the Dex-cAMP treatment (positive control) was defined as 1, any expression level less than 1 from the treatment indicated the inhibition of the mRNA expression of PEPCK.

## 2.8. Glucose production assay

The glucose production assay was performed as previously published with some modifications [9]. Briefly, H4IIE cells treated with Dex-cAMP were added 10 nM insulin, plant extracts, sub-fractions or purified compounds for 5 h at 37 °C in

5% CO<sub>2</sub>. The medium was removed, and the cells were rinsed with phosphate-buffered saline. Glucose production buffer (glucose-free Dulbecco's modified essential medium, pH 7.4, containing 20 mM sodium lactate and 2 nM sodium pyruvate without phenol red) supplemented with dexamethasone and 0.1 mM 8-CPT-cAMP in the presence of 10 nM insulin (Sigma), and plant extracts, sub-fractions, or purified compounds were added, and cells were incubated for an additional 3 h at 37 °C in 5% CO<sub>2</sub>. At the end of the incubation, the glucose concentration was measured from 0.5 mL of culture medium using the Amplex Red glucose assay kit (Invitrogen). The protein concentration quantitated by the Bradford method was used for correction of cell number.

## 2.9. Animal experiments

All animal experiments were performed according to procedures approved by the Rutgers Institutional Animal Care and Use Committee (protocol #04-023). Ten-week-old male C57BL/6J mice were purchased from the Jackson Laboratory (Bar Harbor, ME) and maintained on a high-fat diet containing 60% fat-derived calories (D12492, Research Diets, New Brunswick, NJ) with 12-hour light and dark cycles. The animals were randomized into 3 groups of 10 animals each. The control group was oral gavaged daily with the vehicle solution alone (5% DMSO) and two treatment groups were oral gavaged with 200 mg/kg of the appropriate treatment: crude extract (L) and EtOAc fraction (LE). The animals were weighed weekly for the duration of the experiment. Plasma glucose levels were measured at week 0 and 4 in sub-mandibular vein blood samples using a glucometer (Accu-Chek Advantage, Roche Diagnostics, Indianapolis, IN). A 6 h fast was necessary to allow blood glucose concentrations to arrive at the basal level, plasma glucose concentrations were measured immediately before as 0 h and at 3, 6 h following the treatments of metformin, L, and LE. The oral glucose tolerance test (OGTT) was performed at week 7. The mice were fasted overnight (around 16 h) followed by oral gavage with 2.0 g/kg glucose solution. Glucose levels were subsequently measured using a glucometer at 0, 40, 80, and 130 min after glucose administration.

## 2.10. Data presentation and statistical analysis

Statistical analyses of the experimental observations, expressed as means  $\pm$  SEM, can be assumed to be one-way ANOVA followed with a Tukey's multiple means comparison test, unless otherwise indicated. Treatments were considered significantly different if  $P < 0.05$ .

# 3. Results and discussion

## 3.1. Structure elucidation

Polyphenol compounds exhibit a wide range of pharmacological activities such as anti-cancer, anti-inflammation, and anti-diabetes [22–25]. In our study, ABTS scavenge assay was applied for *in vitro* bio-guided screening based on previous research [26]. Ethyl acetate (LE) fraction was found to exhibit superior activity compared to others (Supplementary Material

S1). Subsequent chemistry analysis indicated that different polyphenol compounds exist in the ethyl acetate fraction (LE) of kinkéliba leaves (Fig. 1). The structure of each compound was elucidated by various spectrometric methods including UV, MS and NMR spectroscopy, or compared with commercial available standards on HPLC. Four flavan compounds were identified as (-)-epicatechin, (-)-epigallocatechin, 3', 4', 5', 5', 7-pentahydroxyflavan and 3', 4', 5', 7-tetrahydroxyflavan using NMR, and the results correlate very well with previous literature reports (Supplementary Material S6) [27,28]. They share identical fragment ions ( $^{1,3}A^+$ ,  $m/z = 139$ ) on mass spectra, which further confirms their flavan identities (Fig. 2A). The chiral centers of compounds epicatechin, epigallocatechin were unambiguously elucidated through the comparison with authentic standards on HPLC (Supplemental Material). This is the first report that these flavan compounds are naturally occurring in kinkéliba. Except the flavans, a number of flavone C-glycosides were separated, characterized and identified. Compound 5, vitexin was unambiguously identified based on UV and MS data and further comparison with authentic standard on HPLC. Compound 6, 7, 8 were tentatively identified as isovitexin, orientin, homoorientin based on their UV and MS spectra along with literature reports. They have identical UV absorption peaks at ~216, 269, 338 nm and characteristic fragment ions  $^{0,2}X^+$  ( $m/z = 313$  or  $329$ ) and  $^{0,3}X^+$  ( $m/z = 359$ ) ions of C-glycosylflavones (Fig. 2B) [29,30]. The O-galloyl derivatives (10–11) of C-glycosylflavones were also separated on HPLC and identified using MS. The structures of 10 and 11 were further confirmed using NMR. The galloyl derivatives of C-glycosylflavones also share identical UV

absorption peaks at ~216, 269, 338 nm [30]. The mass fragment ion  $^{0,2}X^+$  ( $m/z = 481$  or  $465$ ) formed inside the glucose ring and  $[M\text{-galloyl}+H]^+$  ions ( $m/z = 433$  or  $449$ ) further confirms that galloyl group is attached at the 2'' position (Fig. 2C) [31]. Fragment ion ( $m/z = 313, 329$ ) from both  $^{0,2}X^+$  cleavage and galloyl group removal was also observed on the spectra. These O-galloyl derivatives are revealed in kinkéliba for the first time. One known flavonol compound myricetin-3-O-glucoside (9) was also purified and confirmed using NMR [32]. The mass spectra of C-glycosylflavones and their O-galloyl derivatives are illustrated in S4 (Supplementary Material).

### 3.2. Inhibition of PEPCK mRNA expression

The ethanol crude extract (L), ethyl acetate fraction (LE), and its sub-fractions were tested for inhibitory activity of Dex-cAMP induced PEPCK mRNA expression on H4IIE hepatoma cells line. The synthetic compound glucocorticoid dexamethasone and cAMP analog 8-CTP-cAMP were used to up-regulate the expression of PEPCK gene. After the treatment of insulin or tested compounds on H4IIE cell line, the decreased expression levels of PEPCK indicated their inhibitory activity of gluconeogenesis and potential use for diabetes treatment. Cells without any treatments were used as control group and to indicate the basal level of PEPCK expression.  $\beta$ -Actin was used as an internal standard in this assay since its expression won't be affected after the treatments. The crude extract would slightly inhibit the expression of PEPCK at low concentration, and around 16% inhibition was observed at the concentration of 400  $\mu\text{g/ml}$  (Fig. 3A). The ethyl acetate fraction





Fig. 1 – The flavonoids isolated from ethyl acetate fraction (LE) of kinkéliba leaf extract.




**Fig. 2 – The fragmentation pathway of some representative compounds detected in mass spectra. A: (-)-epicatechin, B: orientin, C: 2''-O-galloylorientin.**

further increased the inhibitory activity and dose dependent manner was observed at concentration above 75  $\mu\text{g/ml}$ . The inhibition of LE at 125  $\mu\text{g/ml}$  was up to around 60% (Fig. 3A).

To further identify the main components responsible for the suppression of the mRNA expression of PEPCK, the ethyl acetate extract was further separated into different fractions. Their inhibitory effect and the main compounds contained in each fraction are illustrated in Fig. 3B. The bioactivity of different sub-fractions varies and the ones containing compounds 1–3 and 10–13 showed the highest inhibitory activity (around 83% and 76% inhibition, respectively). The fraction

containing compounds 1–3 (100  $\mu\text{g/ml}$ ) showed stronger inhibitory effect than insulin (10 nM). The major components contained in sub-fraction 1–3 are (-)-epigallocatechin, (-)-epicatechin, and 3', 4', 5', 5, 7-pentahydroxyflavan. Further isolation of each single component for bioassay indicated that compound (-)-epicatechin (2) possessed the best inhibitory activity compared to compound 1, 3, 4, with 32% inhibition at 20  $\mu\text{M}$ . The mixture of different flavones (100  $\mu\text{g/ml}$ ) including 2''-O-galloylvitexin (10), 2''-O-galloylisovitexin (11), 2''-O-galloylorientin (12), 2''-O-galloylhomoorientin (13) also showed potent inhibition activity, even better than insulin of 10 nM.



**Fig. 3** – *In vitro* assays performed on H4IIE rat hepatoma cell line. **A:** The inhibition of the PEPCK mRNA expression of the kinkélíba crude extract (L) and ethyl acetate fraction (LE); **B:** The inhibition of the PEPCK mRNA expression of sub-fractions or compounds isolated from ethyl acetate fraction (LE) of kinkélíba leaves; **C:** The glucose production inhibition of sub-fractions or compounds isolated from ethyl acetate fraction (LE) of kinkélíba leaves.

Sub-fractions 10–11 and 12–13 have relatively lower inhibitory activities compared to 10–13, which is a combination of 10–11 and 12–13. These results suggest that sub-fractions 10–11 and 12–13 may present the inhibitory activity through two or more different mechanistic pathways. While each sub-fraction did not strongly inhibit the expression of PEPCK mRNA, when combined their inhibitory activity increased significantly, suggesting that these compounds may have synergistic effects on the inhibition of PEPCK mRNA expression through two or more different mechanistic pathways.

### 3.3. Inhibition of glucose production


To further assess the effect of down-regulation of PEPCK gene expression on glucose production *in vitro*, compounds pyruvate and lactate, which could be used by H4IIE hepatoma cells for gluconeogenesis, were added to the cell medium. Phytochemicals in the extract, which inhibited the mRNA expression of PEPCK (Fig. 3B), would also inhibit the hepatic gluconeogenesis and result in lower glucose production in cell lines. As shown in Fig. 3C, The fraction containing mainly compound (-)-epicatechin (2), which exhibited the most potent activity in inhibiting the mRNA expression of PEPCK (Fig. 3B), also showed the best inhibitory activity in the glucose production (43% inhibition). Although the mRNA expression

of PEPCK was not decreased by the treatment of compound 3', 4', 5', 7- tetrahydroxyflavan (4) (Fig. 3B), its glucose production was slightly suppressed (16% inhibition) in the hepatoma cell line (Fig. 3C). This discrepancy suggested that other molecular mechanisms may be involved in the anti-diabetic effect of kinkélíba extracts and further investigations are needed.

### 3.4. Animal experiments

#### 3.4.1. Body weight and tissue weight

The daily treatments of kinkélíba samples (L, LE) did not show a significant effect on the body weight change of the animals over the six weeks. The treatment groups have similar weight gain with the control group at all time points we measured, though only week 0 and week 6 are included (Supplementary Material). At week 6, mice fed with LE extract has slight lower body weight and tissue weight compared with the control group indicating kinkélíba could slightly slow down the weight gain, which to some extent correlates with its traditional use for weight loss. This lack of significant weight difference also suggested that the decrease in plasma glucose level is not a result of decreased weight, but rather a therapeutic effect of treatment on glucose production. Similar trend was also observed on the tissue weight, which further supported the non-toxicity of kinkélíba and irrelevance between body/tissue weight and plasma glucose levels.



**Fig. 4** – The results of *in vivo* experiments performed on C57BL/6J mice. (A) Acute glucose experiments; (B) Glucose tolerance test; The data is illustrated as means  $\pm$  SEM, a two-way ANOVA followed by Bonferroni post test was performed with  $P < 0.05$  and  $P < 0.001$  indicated.

#### 3.4.2. *In vivo* plasma glucose levels suppression

The acute blood glucose levels were determined at week 0 and 4. Metformin, an anti-diabetic drug for the treatment of type 2 diabetes, is used as a positive control [33]. As shown in Fig. 4A, after the treatment of leaf extracts, the blood glucose concentration gradually decreased and is comparatively lower than the control group. The ethyl acetate group is slightly more potent than the crude extract based on its higher concentration of polyphenol compounds mentioned above. The blood glucose level decreases by 13.4% at 3 h and 26.1% at 6 h after the treatment of ethyl acetate fraction. Even if the plant extract is not as potent as anti-diabetic drug molecule, metformin, its therapeutic effects on glucose lowering would also be explored in the future and applied as nutraceutical supplements.

#### 3.4.3. Glucose tolerance test

The oral glucose tolerance test was performed after six weeks' administration of kinkéliba leaf extracts. Administration of 2 g/kg glucose to C57BL/6J mice increased the serum glucose from  $159 \pm 35$  mg/dl to  $348 \pm 66$  mg/dl at 40 min,  $356 \pm 84$  mg/dl at 80 min, and returned back to  $243 \pm 13$  mg/dl at 130 min (Fig. 4B). The administration of crude extract (L, 200 mg/kg) suppressed the elevation of serum glucose levels, with glucose level of  $336 \pm 50$  mg/dl and  $315 \pm 62$  mg/dl at 40 min and 80 min, respectively. As expected, the ethyl acetate extract (LE) with concentrated polyphenol compounds showed stronger inhibition of the glucose level upon glucose intake than crude extract. As shown in Fig. 4B, the blood glucose concentration was  $311 \pm 69$  mg/dl and  $297 \pm 58$  mg/dl at 40 and 80 min, respectively. At 80 min, the glucose level of mice treated with LE was 14.5% lower compared with the control group, another indicator of improved glucose metabolism. The results clearly suggested that the kinkéliba could slow down the increase of glucose levels.

polyphenols, contained in African tea plant kinkéliba [34]. Under bio-guided screening, this investigation led to the successful identification of thirteen polyphenols, in which (-)-epigallocatechin, (-)-epicatechin, 3', 4', 5', 5, 7-pentahydroxyflavan, and 3', 4', 5', 7-tetrahydroxyflavan were discovered for the first time in this plant species. Further biological activity studies on purified fractions, which contain different polyphenol compounds, present more in-depth knowledge about the specific functions of polyphenol components. An anti-diabetic activity of kinkéliba was identified and validated *in vitro* through its inhibition of PEPCK mRNA expression and glucose production in H4IIE rat hepatoma cell line. The ethyl acetate sub-fractions containing compounds 1–3, 10–13 showed impressive inhibition of PEPCK mRNA expression and underlying synergistic effects may be involved. Compound 2 (-)-epicatechin has the best *in vitro* PEPCK gene inhibition activity and glucose production inhibition activity. Except PEPCK, other key factors in the development of diabetes, such as Phospho-PKA, phospho-CREB, and PGC-1, should be also further investigated in the future [35–37]. As reflected by the consistence of body weight and tissue weight in *in vivo* experiment, the kinkéliba has no obvious toxicity. Administration of kinkéliba extract, especially the ethyl acetate extract, could decrease the blood glucose levels *in vivo*. At the end of six weeks' experiment, the mice administrated with kinkéliba also showed better glucose tolerance compared with the control group. All these results clearly point to noteworthy *in vitro* and *in vivo* anti-diabetic activity. This is the first report of glucose-lowering activity by kinkéliba herbal tea (*C. micranthum*) and should be explored in further detail, especially the underlying therapeutic mechanisms. Based on its demonstrated bioactivities and the high-demand for the prevention or treatment of diabetes, kinkéliba would also be of interest as nutraceutical supplement in the future and as an herbal tea and in other beverage and food products with this data expanding upon its traditional applications.

## 4. Conclusions

Compared with previous works, this study comprehensively investigated the major phytochemicals, especially the

## Conflicts of interest

All contributing authors have no conflicts of interest to declare.



## Acknowledgements

We thank Babou Diouf and Diatta Malainy, Agri-Business in Sustainable Natural African Plant Product, ASNAPP-Senegal ([www.asnapp.org](http://www.asnapp.org)) with whom we worked to commercialize this indigenous Senegalese plant as an herbal tea as an income generating activity with and for the Senegalese communities. We thank the Senegalese communities who both worked so closely with us and who allowed us to carry out this work with the kinkeliba that was wild harvested on their lands and their support of us to pursue new scientific-driven food and medicinal applications on this species to generate greater market interest in the use of this indigenous Senegalese plant. We thank the USAID for their support of our projects in sub-Saharan Africa to strengthen in-country African natural products industry through their support of the Partnership for Food and Industry in Natural Products (PFID/NP), a university led program supported by the Office of Economic Growth, Agriculture and Trade (EGAT/AG) of the USAID (Contract Award No. AEG-A-00-04-00012-00). We thank the New Jersey Agricultural Experiment Station and the School of Environmental and Biological Sciences at Rutgers and the New Use Agriculture and National Plant Products Program. This work was conducted as part of the dissertation research of Cara Welch and also of Jing Zhen.

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.jfda.2017.05.009>.

## REFERENCES

- [1] Iwu MM. Handbook of African medicinal plants. CRC Press; 2014.
- [2] Juliani HR, Koelliker Y, Bucuk M, Welch C, Tepper BJ, Jefthas E, et al., editors. Quality and consumer studies in the USA of African herbal teas for the natural product industry development in Sub-Sahara Africa. ACS symposium series. Oxford University Press; 2009.
- [3] Le Grand A. Anti-infective phytotherapies of the tree-savannah, Senegal (occidental Africa). III: a review of phytochemical substances and the antimicrobial activity of 43 species. *J Ethnopharmacol* 1989;25:315–38.
- [4] Le Grand A, Wondergem P. Les phytotherapies anti-infectieuses de la foretsavane, Senegal (afrique occidentale) I. Un inventaire. *J Ethnopharmacol* 1987;21:109–25.
- [5] D'Agostino M, Biagi C, Feo Vd, Zollo F, Pizza G. Flavonoids of *Combretum micranthum*. *Fitoterapia* 1990;61.
- [6] Bassene E, Laurance A, Olschwang D, Pousset J. Plantes médicinales africaines: XIX. Dosage de la vitexine par chromatographie liquide haute performance dans un extrait brut de *Combretum micranthum* G. Don. *J Chromatogr A* 1985;346:428–30.
- [7] Bassène E, Olschwang D, Pousset J. Plantes médicinales africaines I: caractérisation de l'inositol et du sorbitol, principes actifs probables du Kinkeliba ("*Combretum micranthum*" G. Don.). *Dakar Med* 1981;26:219–25.
- [8] Simon J, Wu Q, Welch C. Novel piperidine-flavan alkaloid compounds derived from african herb tea kinkeliba as anti-diabetic agents. Patent No. WO/2011/140066., 2011.
- [9] Govorko D, Logendra S, Wang Y, Esposito D, Komarnytsky S, Ribnicky D, et al. Polyphenolic compounds from *Artemisia dracunculus* L. inhibit PEPCK gene expression and gluconeogenesis in an H4IIE hepatoma cell line. *Am J Physiol Endocrinol Metab* 2007;293:E1503–10.
- [10] Collins S, Martin TL, Surwit RS, Robidoux J. Genetic vulnerability to diet-induced obesity in the C57BL/6J mouse: physiological and molecular characteristics. *Physiol Behav* 2004;81:243–8.
- [11] Control CfD, Prevention. National diabetes statistics report: estimates of diabetes and its burden in the United States, 2014. Atlanta, GA: US Department of Health and Human Services; 2014.
- [12] Hamman RF. Genetic and environmental determinants of non-insulin-dependent diabetes mellitus (NIDDM). *Diabetes metab Rev* 1992;8:287–338.
- [13] Roden M, Petersen KF, Shulman GI. Nuclear magnetic resonance studies of hepatic glucose metabolism in humans. *Recent Prog Horm Res* 2001;56:219–37.
- [14] Sun Y, Liu S, Ferguson S, Wang L, Klepcyk P, Yun JS, et al. Phosphoenolpyruvate carboxykinase overexpression selectively attenuates insulin signaling and hepatic insulin sensitivity in transgenic mice. *J Biol Chem* 2002;277:23301–7.
- [15] Tilg H, Moschen AR. Insulin resistance, inflammation, and non-alcoholic fatty liver disease. *Trends Endocrinol Metab* 2008;19:371–9.
- [16] Andersen OM, Markham KR. Flavonoids: chemistry, biochemistry and applications. CRC Press; 2005.
- [17] Babujanarthanam R, Kavitha P, Pandian MR. Quercitrin, a bioflavonoid improves glucose homeostasis in streptozotocin-induced diabetic tissues by altering glycolytic and gluconeogenic enzymes. *Fundam Clin Pharmacol* 2010;24:357–64.
- [18] Guo Y, Su Z-Y, Kong A-NT. Current perspectives on epigenetic modifications by dietary chemopreventive and herbal phytochemicals. *Curr Pharmacol Rep* 2015;1:245–57.
- [19] Hannemann J, Baumann K. Cisplatin-induced lipid peroxidation and decrease of gluconeogenesis in rat kidney cortex: different effects of antioxidants and radical scavengers. *Toxicology* 1988;51:119–32.
- [20] Prince P, Kamalakkannan N. Rutin improves glucose homeostasis in streptozotocin diabetic tissues by altering glycolytic and gluconeogenic enzymes. *J Biochem Mol Toxicol* 2006;20:96–102.
- [21] Singh J, Kakkar P. Antihyperglycemic and antioxidant effect of *Berberis aristata* root extract and its role in regulating carbohydrate metabolism in diabetic rats. *J Ethnopharmacol* 2009;123:22–6.
- [22] Saw CLL, Guo Y, Yang AY, Paredes-Gonzalez X, Ramirez C, Pung D, et al. The berry constituents quercetin, kaempferol, and pterostilbene synergistically attenuate reactive oxygen species: involvement of the Nrf2-ARE signaling pathway. *Food Chem Toxicol* 2014;72:303–11.
- [23] Guo Y, Shu L, Zhang C, Su Z-Y, Kong A-NT. Curcumin inhibits anchorage-independent growth of HT29 human colon cancer cells by targeting epigenetic restoration of the tumor suppressor gene DLEC1. *Biochem Pharmacol* 2015;94(2):69–78.
- [24] Zhen J, Guo Y, Villani T, Carr S, Brendler T, Mumbengegwi DR, et al. Phytochemical analysis and anti-inflammatory activity of the extracts of the African medicinal plant *Ximenia caffra*. *J Anal Methods Chem*. 2015;2015(4):1–9.
- [25] Anhê FF, Desjardins Y, Pilon G, Dudonné S, Genovese MI, Lajolo FM, et al. Polyphenols and type 2 diabetes: a prospective review. *PharmaNutrition* 2013;1:105–14.

- [26] Zhen J, Villani TS, Guo Y, Qi Y, Chin K, Pan M-H, et al. Phytochemistry, antioxidant capacity, total phenolic content and anti-inflammatory activity of *Hibiscus sabdariffa* leaves. *Food Chem* 2016;190:673–80.
- [27] Davis AL, Cai Y, Davies AP, Lewis JR. H-1 and C-13 NMR assignments of some green tea polyphenols. *Magn Reson Chem* 1996;34:887–90 [In English].
- [28] Teixeira AF, Alcântara AFdC, Piló-Veloso D. Structure determination by <sup>1</sup>H and <sup>13</sup>C NMR of a new flavan isolated from *Brosimum acutifolium*: 4', 7-dihydroxy-8-prenylflavan. *Magn Reson Chem* 2000;38:301–4.
- [29] Peng X, Zheng Z, Cheng K-W, Shan F, Ren G-X, Chen F, et al. Inhibitory effect of mung bean extract and its constituents vitexin and isovitexin on the formation of advanced glycation endproducts. *Food Chem* 2008;106:475–81.
- [30] Latté KP, Ferreira D, Venkatraman M, Kolodziej H. O-Galloyl-C-glycosylflavones from *Pelargonium reniforme*. *Phytochemistry* 2002;59:419–24.
- [31] Cabrera GM. Mass spectrometry in the structural elucidation of natural products: glycosides. *Phytochem: Adv Res Res* 2006:1–22. Signpost, Kerala.
- [32] Castillo-Munõz N, Gomez-Alonso S, Garcia-Romero E, Gómez MV, Velders AH, Hermosin-Gutierrez I. Flavonol 3-O-glycosides series of *Vitis vinifera* cv. Petit Verdot red wine grapes. *J Agric Food Chem* 2008;57:209–19.
- [33] Bailey CJ, Turner RC. Metformin. *N. Engl J Med* 1996;334:574.
- [34] Chika A, Bello SO. Antihyperglycaemic activity of aqueous leaf extract of *Combretum micranthum* (Combretaceae) in normal and alloxan-induced diabetic rats. *J Ethnopharmacol* 2010;129:34–7.
- [35] Wang Y, Li G, Goode J, Paz JC, Ouyang K, Screatton R, et al. Inositol-1, 4, 5-trisphosphate receptor regulates hepatic gluconeogenesis in fasting and diabetes. *Nature* 2012;485:128–32.
- [36] Koo S-H, Flechner L, Qi L, Zhang X, Screatton RA, Jeffries S, et al. The CREB coactivator TORC2 is a key regulator of fasting glucose metabolism. *Nature* 2005;437:1109–11.
- [37] Yoon JC, Puigserver P, Chen G, Donovan J, Wu Z, Rhee J, et al. Control of hepatic gluconeogenesis through the transcriptional coactivator PGC-1. *Nature* 2001;413:131–8.