

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect


journal homepage: [www.jfda-online.com](http://www.jfda-online.com)

## Corrigendum

# Corrigendum to “High-throughput liquid chromatography tandem mass spectrometry method for simultaneous determination of fampridine, paroxetine, and quinidine in rat plasma: Application to *in vivo* perfusion study” [J Food Drug Anal 24 (2016) 866–875]

Suneetha Achanti <sup>a,\*</sup>, Raja Rajeswari Katta <sup>b</sup><sup>a</sup> Department of Pharmaceutical Analysis, Hindu College of Pharmacy, Guntur, Andhra Pradesh, India<sup>b</sup> College of Pharmaceutical Sciences, Acharya Nagarjuna University, Guntur, Andhra Pradesh, India

The GA (Graphical Abstract) is corrected to include the 2 missing points. One is the Heating mat is drawn for the anaesthetized rat and secondly, the perfusion solution was used in red color due to phenol red which is actually done at the time of experiment and collected the perfusate in eppendorf tubes only.

DOI of original article: <http://dx.doi.org/10.1016/j.jfda.2016.03.004>.

\* Corresponding author. Department of Pharmaceutical Analysis, Hindu College of Pharmacy, Amaravathi Road, Guntur 522 002, Andhra Pradesh, India.

E-mail address: [raj.pharm@rediffmail.com](mailto:raj.pharm@rediffmail.com) (S. Achanti).<http://dx.doi.org/10.1016/j.jfda.2017.01.006>

1021-9498/Copyright © 2017, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).