

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.jfda-online.com

Original Article

Antioxidant activity, neuroprotective properties and bioactive constituents analysis of varying polarity extracts from *Eucalyptus globulus* leaves



Elena González-Burgos ^{a,*}, Mindaugas Liaudanskas ^{b,c,1}, Jonas Viškelis ^c, Vaidotas Žvikas ^d, Valdimaras Janulis ^d, M. Pilar Gómez-Serranillos ^a

^a Department of Pharmacology, Pharmacognosy and Botany, Faculty of Pharmacy, University Complutense Madrid, Madrid, Spain

^b Institute of Pharmaceutical Technologies of the Faculty of Pharmacy of Lithuanian University of Health Sciences, Kaunas, Lithuania

^c Institute of Horticulture, Lithuanian Research Centre for Agriculture and Forestry, Kaunas, Lithuania

^d Department of Pharmacognosy, Faculty of Pharmacy, Lithuanian University of Health Sciences, Kaunas, Lithuania

ARTICLE INFO

Article history:

Received 22 February 2018

Received in revised form

11 May 2018

Accepted 22 May 2018

Available online 18 June 2018

Keywords:

Eucalyptus globulus

Neuroprotection

Antioxidants

Bioactive constituents

ABSTRACT

Eucalyptus globulus is employed as herbal tea and therapeutical purposes. In this work, it is investigated for first time the neuroprotective activities, based on antioxidant properties, of varying polarity extracts (acetone, ethanol and methanol) from *E. globulus* leaves and elucidate their main bioactive constituents. Methanol and acetone extracts contained the highest phenolic compounds amount and chlorogenic acid was the major compound identified by UPLC-ESI-MS/MS. Moreover, the three tested extracts showed significant antioxidant properties, varying their potency depending on the *in vitro* technique used. Furthermore, *E. globulus* extracts were effective in ameliorating H₂O₂-induced oxidative stress by increasing cell viability, GSH levels and antioxidant enzymes activity and, by decreasing ROS production and lipid peroxidation levels in SH-SY5Y cells. Taken together, *E. globulus* leaves extracts could be used as raw material for food and pharmaceutical supplements for their high content in antioxidant compounds with health benefits properties against oxidative stress.

Copyright © 2018, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Eucalyptus globulus Labill. (Lan An) is an evergreen tree native to Australia and extensively cultivated in Mediterranean and subtropical areas for firewood and pulp [1]. *E. globulus* leaves

are traditionally employed for asthma and bronchitis treatment. Moreover, they are used to make herbal tea. Furthermore, recent studies have demonstrated antimicrobial, antifungal, anthelmintic and anti-diabetic properties for leaves extracts and essential oils [2].

* Corresponding author.

E-mail address: elenagon@ucm.es (E. González-Burgos).

¹ These authors contributed equally to this work.

<https://doi.org/10.1016/j.jfda.2018.05.010>

1021-9498/Copyright © 2018, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Oxidative stress is the result of a cell redox imbalance, leading to biomolecules structure and function disturbance. The brain is vulnerable to reactive oxygen species (ROS) injury because of its high concentrations of polyunsaturated fatty acids, high oxygen consumption and lack of antioxidant defense mechanisms. Particularly, neurons are especially susceptible to oxidative damage owing to its reduced glutathione low levels and its minimal regenerative capacity. Extensive evidence points to oxidative stress as major contributor in the pathogenesis of Alzheimer's disease (AD) and Parkinson's disease (PD). Central nervous system pathologies account for 12% of all deaths worldwide and this number continues to rise due to aging-associated increased incidence [3].

Exogenous antioxidants, which prevent from redox imbalance by scavenging ROS, chelating metals and up-regulating enzymatic and non-enzymatic system, constitute the most promising therapeutic strategy for oxidative stress-mediated pathologies. Hence, due to the increasing number of people suffering from neurodegenerative diseases and the lack of effective treatments, there is a serious need to find new antioxidant sources that slow CNS pathologies progression. Polyphenols are presently getting great attention for their antioxidant properties and their potential role as neuro-protective agents. They act as effective electron and hydrogen donors, singlet oxygen quenchers and reducing agents, preventing from ROS harmful effects [4].

The aim of this study was to investigate for first time the neuroprotective effect, based on its antioxidant properties, of acetone, ethanol and methanol extracts of *E. globulus* leaves against H₂O₂-induced oxidative stress in neuroblastoma cells.

2. Methods

2.1. Reagents

Acetonitrile, 2,2'-azinobis(3-ethylbenzothiazoline-6-sulphonic acid (ABTS), 3-(4,5-Dimethyl-2-thiazolyl)-2,5-diphenyltetrazolium bromide (MTT), 2,7-dichloro-dihydro-fluorescein diacetate (DCFH-DA), dimethyl sulphoxide (DMSO), 2,2-diphenyl-1-picrylhydrazyl (DPPH), Dulbecco's modified Eagle's medium (DMEM), hydrogen peroxide and 2,4,6-triphenyl-1,3,5-triazine (TPTZ) were acquired from Sigma–Aldrich (St. Louis, MO, USA). All standards were also purchased from Sigma–Aldrich (St. Louis, MO, USA) except for luteolin-4-glucoside and isorhamnetin 3-O-glucoside which were from Extrasynthese (Genay, France). Fetal bovine serum (FBS), phosphate-buffered saline (PBS) and gentamicin were from Gibco (Invitrogen, Paisley, UK).

2.2. Plant material

E. globulus leaves (manufacturer JSC “Acorus Calamus”, Švenčionys, Lithuania) were ground to a powder using a mill (IKA® A11 basic, Staufen im Breisgau, Germany). Loss on drying before analysis was determined by drying about 1 g of milled leaves in a moisture analyzer (Precisa HA 300, Precisa Instruments AG, Dietikon, Switzerland) to complete evaporation of water and volatile compounds (drying temperature: 105 °C).

2.3. Extraction

Raw material (125 g) was soaked for 3 h in 70% (v/v) methanol, ethanol or acetone. Soaked raw material was transferred to a percolator, covered with extractant (250 mL) and left to macerate (48 h). Then it was percolated (rate 0.3 mL min⁻¹) and high concentration extract (85% of total extract amount) was obtained. Low concentration extract was decanted and it was evaporated using rotary evaporator (IKA® HB 10, Staufen im Breisgau, Germany) up to 15% of total liquid extract amount. Remaining part of low concentration extract was transferred to a single container with high concentration extract. Organic phase of liquid extract was evaporated using rotary evaporator and remaining aqueous phase was lyophilised using lyophilisator Zirbus (Zirbus technology GmbH, Germany) at 0.01 mbar pressure and, condenser temperature of –85 °C.

2.4. UPLC-ESI-MS/MS conditions

Separation of phenolic compounds was performed with Acquity H-class UPLC system (Waters, USA) equipped with triple quadrupole tandem mass spectrometer (Xevo, Waters, USA) with an electrospray ionization source (ESI) to obtain MS/MS data. YMC Triart C18 (100 × 2.0 mm; 1.9 µm) column (YMC Europe GmbH, Dislanken, Germany) was used for analysis. Column temperature was maintained at 40 °C. Gradient elution was performed with mobile phase consisting of 0.1% formic acid water solution (solvent A) and acetonitrile (solvent B) with flow rate set to 0.5 mL min⁻¹. Linear gradient profile was applied as follows for solvent A: initially 95% for 1 min; to 70% over 4 min; 50% over 7 min; 95% over 2 min. Negative electrospray ionization was applied for analysis: capillary voltage –2 kV, source temperature 150 °C, desolvation temperature 400 °C, desolvation gas flow 700 L h⁻¹, cone gas flow 20 L h⁻¹. Collision energy and cone voltage were optimized for each compound separately.

2.5. Validation of the method

The validation of the UPLC-ESI-MS/MS method was performed on the basis of the US Food and Drug Administration [5].

The evaluation of the selectivity of the method for peak identification and purity were based on the comparison of the retention times and MS spectra of the analytes with those of the standard compounds.

The limit of detection (LOD) and limit of quantification (LOQ) of the analytes were assessed by comparing the peak height to the baseline noise. The signal-to-noise ratio was 3:1 for a limit of detection and it was 10:1 for a limit of quantification. The determined LOD varied from 0.4 ng/mL to 32.51 ng/mL and the determined LOQ varied from 1.2 ng/mL to 100.25 ng/mL. Calibration curves were obtained by plotting the peak areas of analytical standards. The estimated determination coefficients (R²) of calibration curves ≥0.96 (Supporting information).

The precision and accuracy were analyzed by evaluating quality control standard mixture at twice the level of LOQ. Accuracy and precision were determined by analyzing samples on three consecutive days (6 × 3 injections). Accuracy and

precision were estimated using relative standard deviation (RSD). The calculated % RSD for the peak area varied from 2.7% (p-coumaric acid) to 9.16% ((-)-Epicatechin gallate). The mean values of accuracy and precision were below acceptable 15% at two times the LLOQ level.

2.6. Antioxidant activity evaluation

DPPH method. DPPH solution in 96.3% v/v ethanol (3 mL, 6×10^{-5} M) was mixed with 10 μ L of extracts. A decrease in absorbance was determined at $\lambda = 515$ nm in a double beam UV/VIS spectrophotometer M550 (Spectronic Camspec, Garforth, UK) [6]. This spectrophotometer was employed for all antioxidant experiments. Results are expressed as mmol Trolox equivalent (TE) per gram of absolutely dry weight (DW).

ABTS \cdot + method. 3 mL of ABTS \cdot + solution was mixed with 10 μ L of extracts. A decrease in absorbance was measured at $\lambda = 734$ nm [7].

TFPH \cdot + method. 3 mL of TFPH \cdot + solution was mixed with 10 μ L of extracts and absorbance was measured at $\lambda = 502$ nm [8].

FRAP Assay. FRAP solution included TPTZ (0.01 M dissolved in 0.04 M HCl), $\text{FeCl}_3 \times 6\text{H}_2\text{O}$ (0.02 M in water), and acetate buffer (0.3 M, pH 3.6) (ratio 1:1:10). 3 mL of a freshly prepared FRAP reagent was mixed with 10 μ L of extracts. An increase in absorbance was recorded at $\lambda = 593$ nm [9].

CUPRAC Assay. CUPRAC solution included copper (II) chloride (0.01 M in water), ammonium acetate buffer solution (0.001 M, pH = 7), and neocuproine (0.0075 M in ethanol) (ratio 1:1:1). 3 mL of CUPRAC reagent was mixed with 10 μ L of extracts. An increase in absorbance was recorded at $\lambda = 450$ nm [10].

2.7. Cell culture

Human neuroblastoma SH-SY5Y cells were grown in DMEM, FBS (10%) and gentamicin (50 mg mL $^{-1}$) at 37 °C in humidified atmosphere of 5% CO $_2$. The SH-SY5Y cell line was provided by the University of Alcalá (Madrid, Spain), CAI Medicine and Biology, Cell Culture Unity.

Extracts were first dissolved in DMSO and then diluted to the desired concentrations in PBS (final concentration of $\leq 0.1\%$ DMSO). Cells were pre-treated with different concentrations (from 5 μ g mL $^{-1}$ to 1000 μ g mL $^{-1}$) prior to H $_2$ O $_2$ (0.1 mM, 30 min).

2.8. MTT cytotoxic assay

MTT dye solution (2 mg mL $^{-1}$) was added and after 1 h of incubation, formazan crystals were dissolved with DMSO. Absorbance was measured spectroscopically at 570 nm (LT-4000, Microplate Reader, Labtech International Ltd, UK). Results were expressed as relative percentages, considering absorbance readings of untreated cells as 100% [11].

2.9. Intracellular ROS production

Cells were incubated with DCFH-DA (10 mM) in PBS-glucose for 30 min in darkness at 37 °C. Fluorescence was recorded

at $\lambda_{\text{exc}} = 480$ nm and at $\lambda_{\text{em}} = 510$ nm in a microplate reader FLx800, Bio-Tek Instrumentation [12].

2.10. Reduced and oxidized glutathione content

To measure GSH concentration, supernatants of cell lysate samples (50 μ L) were incubated with ortho-phthalaldehyde (20 μ L, 1 mg mL $^{-1}$ metanol) for 15 min. To measure GSSG concentration, supernatants of cell lysate samples (50 μ L) were first incubated with N-ethylmaleimide (3 μ L) for 5 min and then with ortho-phthalaldehyde (20 μ L, 1 mg mL $^{-1}$ metanol) for 15 min. Fluorescence values were measured at $\lambda_{\text{exc}} = 350$ nm and $\lambda_{\text{em}} = 420$ nm. Results were calculated using GSH and GSSG calibration curves and expressed as GSH/GSSG ratio [13].

2.11. TBARS assay

Lipid peroxidation was determined by thiobarbituric acid reactive substances (TBARS) assay [14]. Briefly, cellular lysates (50 μ L) were incubated with thiobarbituric acid (TBA), trichloroacetic acid (TCA) and HCl reagent in boiling water bath for 10 min. Then, the mixture was centrifuged (3000 rpm, 10 min, 4 °C). Absorbance was measured at 535 nm in a microplate reader Digiscan 340 (Asys Hitech GmbH, Eugendorf, Austria). Total TBARS content was expressed as % of control.

2.12. Antioxidant enzymes activities

For total lysates, cell pellets were resuspended in a lysis buffer pH 7.4 (Tris 25 mM, EDTA 1 mM NaCl 150 mM, and 0.1% Triton X-100) contained proteinase inhibitors (20 μ L mL $^{-1}$, leupeptin, 35 μ L mL $^{-1}$ PMSF and 10 μ L mL $^{-1}$ pepstatin) for 20 min at 4 °C. Then, cells were centrifuged at 2500 g for 10 min at 4 °C and supernatants were employed for enzymatic assays.

Catalase (CAT) activity. Briefly, total lysates (30 μ L) were dissolved in 150 mM of H $_2$ O $_2$ (670 μ L) dissolved in 50 mM of phosphate buffer (pH 7.4). Hydrogen peroxide decomposition was recorded spectrophotometrically (UVIKON 930, Kontron Instruments, UK) for 1 min at 240 nm [15].

Superoxide dismutase (SOD). Briefly, the assay mixture consisted of 1 mM EDTA (20 μ L), 96 mM NBT (20 μ L), 50 mM hydroxylamine hydrochloride (5 μ L) and cellular lysate (25 μ L). Absorbance was measured at 560 nm using a LT-4000, Microplate Reader, Labtech International Ltd, UK [16].

Glutathione reductase (GR). Briefly, the assay mixture consisted of 20 mM EDTA (20 μ L), 50 mM phosphate buffer (40 μ L, pH 7.4), 2 mg mL $^{-1}$ NADPH (20 μ L), 5 mM GSSG (20 μ L) and cellular lysate (25 μ L). Absorbance was measured at 340 nm using a microplate reader [17].

Glutathione peroxidase (GPx). Briefly, the assay mixture consisted of 50 mM phosphate buffer (100 μ L, pH 7.4), cellular lysate (20 μ L), 0.048 U of GR, 10 mM GSH (20 μ L), 2 mg mL $^{-1}$ NADPH (20 μ L) and 63.5 mM H $_2$ O $_2$ (20 μ L). Absorbance was measured for at 340 nm using a microplate reader [18].

Table 1 – Qualitative and quantitative analysis of bioactive constituents of varying polarity extracts from *Eucalyptus globulus* leaves.

	Compound	m/z	Daughter ion	Cone voltage V	Collision energy eV	Acetone extract C, $\mu\text{g g}^{-1}$ DW	Ethanol extract C, $\mu\text{g g}^{-1}$ DW	Methanol extract C, $\mu\text{g g}^{-1}$ DW
1	3,4-Dihydroxybenzoic acid (Protocatechuic acid)	153	81	32	20	207.88 ± 8.22 ^b	90.59 ± 3.26 ^{a,c}	199.96 ± 6.20 ^b
2	p-coumaric acid	163	93	28	22	34.23 ± 0.92 ^{b,c}	20.10 ± 0.06 ^a	22.05 ± 0.78 ^a
3	Vanillic acid	167	152	26	12	8.13 ± 0.38 ^{b,c}	0.19 ± 0.01 ^{a,c}	6.30 ± 0.27 ^{a,b}
4	Gallic acid	169	51	36	30	415.42 ± 19.32 ^c	334.70 ± 12.36	270.19 ± 10.31 ^a
5	Caffeic acid	179	107	36	22	6.99 ± 0.27 ^c	6.42 ± 0.27 ^c	3.53 ± 0.15 ^{a,b}
6	Quinic acid	191	85	40	26	1801.68 ± 87.32	2159.58 ± 0.93	1994.00 ± 80.36
7	Ferulic acid	193	134	32	18	35.71 ± 1.65 ^b	17.55 ± 0.79 ^{a,c}	41.53 ± 1.26 ^b
8	Sinapic acid	223	208	32	14	2.42 ± 0.08 ^b	ND	0.21 ± 0.01 ^a
9	Chrysin	253	63	56	30	ND	0.54 ± 0.02	0.65 ± 0.02
10	Apigenin	269	117	54	36	25.00 ± 1.13 ^b	459.00 ± 19.32 ^{a,c}	2.79 ± 0.09 ^b
11	Naringenin	271	151	46	18	19.17 ± 0.87 ^b	26.13 ± 1.12 ^{a,c}	14.39 ± 0.65 ^b
12	Phloretin	273	167	42	16	31.72 ± 1.30	36.92 ± 1.56 ^c	25.59 ± 0.16 ^b
13	Luteolin	285	133	58	36	157.21 ± 6.29 ^{b,c}	84.07 ± 3.30 ^a	69.70 ± 2.73 ^a
14	Scutellarein	285	117	58	36	0.77 ± 0.03	ND	1.16 ± 0.04
15	Eriodictyol	287	151	96	14	3.49 ± 0.14 ^c	1.90 ± 0.07 ^c	11.15 ± 0.47 ^{a,b}
16	(-)-Epicatechin	289	123	60	34	1.72 ± 0.05 ^c	ND	12.23 ± 0.57 ^a
17	(+)-Catechin	289	123	60	34	54.34 ± 2.50	ND	70.98 ± 3.46
18	Diosmetin	299	284	46	20	32.24 ± 0.85 ^{b,c}	9.13 ± 0.37 ^a	8.97 ± 0.41 ^a
19	Quercetin	301	151	48	20	685.05 ± 33.47 ^c	828.68 ± 35.63 ^c	420.95 ± 17.36 ^{a,b}
20	Ellagic acid	301	273	64	38	28.06 ± 1.02 ^b	7.73 ± 0.31 ^a	ND
21	Taxifoline	303	151	38	18	ND	1.50 ± 0.06	1.23 ± 0.04
22	Isorhamnetin	315	300	44	22	13.42 ± 0.48 ^c	15.28 ± 0.65 ^c	8.28 ± 0.35 ^{a,b}
23	Chlorogenic acid	353	191	32	14	15,377.21 ± 589.31 ^b	7476.25 ± 303.26 ^{a,c}	18,768.28 ± 803.32 ^b
24	Neochlorogenic acid	353	191	32	14	292.41 ± 26.36 ^c	338.58 ± 15.63 ^c	826.57 ± 30.26 ^{a,b}
25	Rosmarinic acid	359	161	36	16	667.31 ± 29.32 ^{b,c}	30.06 ± 1.38 ^{a,c}	297.86 ± 11.32 ^{a,b}
26	Apigenin 7-glucoside	431	268	66	32	9.42 ± 0.33 ^b	6.39 ± 0.23 ^a	8.15 ± 0.32 ^b
27	Vitexin	431.1	311	50	22	409.17 ± 18.01 ^b	211.77 ± 7.03 ^{a,c}	410.96 ± 18.96 ^b
28	Quercetin 3-O- α -L-arabinofuranoside (Avicularin)	433	301	50	20	131.34 ± 5.64 ^b	51.49 ± 2.01 ^{a,c}	122.37 ± 4.65 ^b
29	Quercetin 3-O- α -L-arabinopyranoside (Guaiajaverin)	433	300	56	26	41.04 ± 1.29	ND	43.60 ± 1.36
30	Phloretin-2'-O-glucoside (Phlorizin)	435	273	42	14	4666.50 ± 203.69 ^b	3382.28 ± 130.60 ^{a,c}	5515.93 ± 252.36 ^b
31	(-)-Epicatechin gallate	441	169	40	16	25.58 ± 0.85	ND	18.63 ± 0.75
32	Homoorientin (Isoorientin)	447	327	50	24	423.09 ± 19.21 ^b	124.80 ± 5.32 ^{a,c}	411.89 ± 19.32 ^b
33	Kaempferol-3-O-glucoside (Astragaln)	447	284	54	28	224.83 ± 8.63 ^b	109.15 ± 4.74 ^{a,c}	213.63 ± 8.36 ^b
34	Luteolin-4-glucoside	447	285	36	16	127.39 ± 5.21	ND	89.57 ± 3.72
35	Luteolin-7-O-glucoside (Cynaroside)	447	285	66	26	8.84 ± 0.32 ^{b,c}	2.88 ± 0.11 ^{a,c}	14.88 ± 0.63 ^{a,b}
36	Orientin	447	327	50	20	407.37 ± 16.30 ^b	106.32 ± 4.96 ^{a,c}	425.90 ± 19.26 ^b
37	Quercetin 3-O-rhamnoside (Quercitrin)	447	300	50	26	268.80 ± 11.23 ^{b,c}	60.93 ± 2.36 ^{a,c}	178.86 ± 7.53 ^{a,b}
38	Kaempferol-3-O-glucuronide	461	285	36	20	6007.94 ± 265.32 ^b	226.87 ± 8.56 ^{a,c}	5543.54 ± 203.69 ^b
39	Quercetin 3-O-glucoside (Isoquercitrin)	463	301	52	28	1143.85 ± 49.63 ^b	101.99 ± 4.03 ^{a,c}	1408.52 ± 69.31 ^b
40	Isorhamnetin-3-O-glucoside	477	314	60	28	39.01 ± 1.82	ND	48.90 ± 2.06
41	Quercetin malonylglycoside	549	300	20	38	3.24 ± 0.14	ND	1.17 ± 0.04

42	Luteolin-7-rutinoside (Scolymoside)	593	285	82	36	14.28 ± 0.63 ^c	ND	1.88 ± 0.07 ^a
43	Quercetin-3-O-rutinoside (Rutin)	609	300	70	38	1986.30 ± 92.35 ^b	30.49 ± 1.32 ^{a,c}	2010.50 ± 87.31 ^b
44	Isorhamnetin-3-O-rutinoside	623	315	70	32	26.06 ± 1.02 ^b	4.57 ± 0.18 ^{a,c}	25.37 ± 0.98 ^b
	Total (µg g ⁻¹ DW)					35,865.62 ± 1456.32 ^b	16,364.81 ± 625.32 ^{a,c}	39,572.78 ± 1563.32 ^b

Statistical significances of quantitative composition are shown in superscripts. a: Statistically significant differences versus acetone extract values. b: versus ethanol extract values. c: versus methanol extract values.

2.13. Statistical analysis

Experiments were performed in triplicate and results were expressed as mean ± SD. One-way ANOVA, followed by Fisher's least-significant-difference (LSD) test ($P \leq 0.05$) was performed using GraphPad Prism v.5 software.

3. Results and discussion

3.1. Evaluation of phenolic compounds using UPLC-ESI-MS/MS

Qualitative and quantitative phenolic composition of *E. globulus* extracts was investigated using UPLC-MS/MS (Table 1). Total phenolic compounds amount in acetone and methanol extracts was similar, being 35,865.6 µg g⁻¹ and 39,572.8 µg g⁻¹, respectively. Total phenol content in ethanol extract (16,364.8 ± 625.3 µg g⁻¹) was 2.2 and 2.4 times less than total amount of these compounds in acetone and methanol extracts, respectively. These results confirm previous works regarding the extraction solvent effect on biological active compounds composition [19].

UPLC-ESI-MS/MS analysis showed that the highest proportion of phenolic compounds in extracts were compounds of flavonoid group (i.e. quercetin, luteolin, kaempferol, isorhamnetin, phloretin). The maximum total amount of flavonoids was determined in methanol extract (17,142.3 ± 702.2 µg g⁻¹) and the lowest in ethanol extract (5883.1 ± 203.6 µg g⁻¹). The largest group of flavonoids was quercetin glycosides with antioxidant, anti-cancer, antimicrobial and cardiovascular protective properties [20].

Other flavonoids in lesser amounts were isorhamnetin and its 3 glycosides, luteolin and its 2 C-glycosides, luteolin-4'-methyl ether, apigenin and its 2 glycosides, 2 kaempferol glycosides, phloretin and phloretin. Particularly, kaempferol-3-O glucoside was predominant in methanol and acetone samples while phlorizin was in ethanol extract (Table 1). Moreover, the monomeric flavan-3-ol compounds (+)-catechin, (-)-epicatechin and (-)-epicatechin gallate were quantified; they have antioxidant, anti-inflammatory, hypocholesterolemic and vasodilatory actions [21].

Phenolic acids including hydroxybenzoic acids (protocatechuic acid and vanillic acid) and hydroxycinnamic acids (caffeic acid, ferulic acid, synapic acid, p-coumaric acid and chlorogenic acid) were also identified. The methanol extracts had the highest phenolic acid content (20,436.5 ± 808.4 µg g⁻¹) whereas the lowest concentration was for ethanol extract (8314.4 ± 352.2 µg g⁻¹). The chlorogenic acid was the most abundant analyte; it has antioxidant, anti-inflammatory, renoprotective, hepatoprotective, anti-diabetic and antilipidemic properties [22].

3.2. Antioxidant activity

Scientific studies have proven that antioxidants have the capacity to slow down PD and AD progression. There is growing interest in antioxidants to pursue opportunities to apply them in medicine, pharmacy, cosmetic and food industry [23].

We then investigated *in vitro* antiradical and reductive activity of acetone, ethanol and methanol *E. globulus* extracts (Table 2). Herbal extracts have a variety of compounds which act as antioxidant through different reaction mechanisms. Therefore, performing a single test of antioxidant activity is not adequate, it is recommended to use at least two different methods [24].

Herbal products can exert their antioxidant action through two main mechanisms: hydrogen atom abstraction (HAT) and single electron transfer (SET). Antioxidant effects depend on chemical antioxidant structure, physico-chemical properties (solubility, partition coefficient) and medium properties (type of solvent, pH) [25]. The *in vitro* antioxidant properties of extracts should be determined using different modelled systems

with different reaction mechanisms with the aim to predict *in vivo* antioxidant activity. Particularly, we assessed 5 different antioxidant capacity assays (ABTS, CUPRAC, DPPH, FRAP, and TFPH). ABTS, DPPH and TFPH assays are based on the ability of antioxidants to scavenge ABTS^+ , DPPH^+ or TFPH^+ free radicals [6–8] whereas FRAP and CUPRAC assays measure the capability of antioxidants to reduce Fe(III) to Fe(II) or Cu(II) to Cu(I) [10]. The knowledge of the antioxidant activity could be useful for the standardization of the raw materials of *E. globulus* leaves extracts.

The present study showed that the highest *in vitro* antioxidant activity by ABTS and CUPRAC methods was for acetone extracts (10.1 and 3.65 mmol g^{-1} , respectively). Moreover, the highest antioxidant activity determined by DPPH method was

Table 2 – Antioxidant activity of extracts of *Eucalyptus globulus* leaves assessed by ABTS, CUPRAC, DPPH, FRAP and TFPH assays.

	ABTS TE, mmol g^{-1} DW	CUPRAC TE, mmol g^{-1} DW	DPPH TE, mmol g^{-1} DW	FRAP TE, mmol g^{-1} DW	TFPH TE, mmol g^{-1} DW
Acetone extract	10.06 ± 0.44^b	$3.65 \pm 0.10^{b,c}$	0.97 ± 0.01^c	1.31 ± 0.19^b	1.13 ± 0.10^b
Ethanol extract	$1.69 \pm 0.07^{a,c}$	$0.48 \pm 0.03^{a,c}$	0.96 ± 0.04^c	$9.79 \pm 0.24^{a,c}$	$1.78 \pm 0.07^{a,c}$
Methanol extract	9.91 ± 0.32^b	$2.69 \pm 0.08^{a,b}$	$1.56 \pm 0.03^{a,b}$	1.10 ± 0.08^b	0.91 ± 0.13^b

Statistical significances of antioxidant capacities values are shown in superscripts letters: a statistically significant differences versus values of acetone extract; b versus values of ethanol extract and c versus values of methanol extract ($p < 0.05$).




Fig. 1 – (A) Effects of *Eucalyptus globulus* extracts on cell viability. Human neuroblastoma SH-SY5Y cells were treated with various concentrations (from $5 \mu\text{g mL}^{-1}$ to $1000 \mu\text{g mL}^{-1}$) of acetone extract, ethanol extract and methanol extract for 24 h. (B) Protective effects of *Eucalyptus globulus* extracts against hydrogen peroxide toxicity. Human neuroblastoma SH-SY5Y cells were pretreated with non-cytotoxic concentrations of acetone extract, ethanol extract and methanol extract for 24 h, prior to the toxic agent hydrogen peroxide (0.1 mM , 30 min). The viable number of cells was determined by using MTT assay. Results were expressed as percentage of untreated cells. Each point represents mean values \pm SEM of three independent experiments. * $p < 0.05$ compared with control and # $p < 0.05$ compared with hydrogen peroxide.

found for methanol extract (1.6 mmol g^{-1}). Finally, the highest values for FRAP and TFPH methods were 9.8 and 1.8 mmol g^{-1} , respectively, in ethanol extracts (Table 2). Such results may have been due to the characteristics of these methods and antioxidants physicochemical properties: different environment pH, hydrophilic–lipophilic properties, unequal different antioxidant distribution between lipophilic and hydrophilic media, differences in reaction mechanisms and in the qualitative and quantitative composition of antioxidant active compounds. The pH of the environment is one of the most affecting factors for antioxidant activity. Since the degree of solubility and stability of different antioxidant forms is highly dependent on the pH of the environment and different human body fluids have different pH, each exogenous antioxidant has a specific antioxidant capacity *in vivo*, which may differ from only one *in vitro* method used. In our study, the presence of low pH (FRAP and TFPH methods), the strongest antioxidant activity was found in ethanol *E. globulus* extracts, while when the pH value was close to the human body fluids (ABTS and CUPRAC methods), the strongest antioxidant activity was found for acetone extracts. Many of phenolic compounds antioxidative activity are higher when reaction medium pH is close to neutral or weakly alkaline [6–10, 26], therefore pH of medium is very important factor deciding biologically active compounds effect potency. Yet different other results were obtained by evaluating the antiradical activity of extracts in non-aqueous organic environment using DPPH method. The strongest antiradical activity was found in methanol extracts, which, when tested by other methods, did not showed such strong antioxidant activity. Such unequal results could derive because of applied methods deficiencies limiting totally adequate and precise evaluation of extracts biologically active compounds antioxidative activity *in vitro*: application of DPPH method for evaluation of hydrophilic antioxidants activity is very complicated [27], FRAP method can not determine antioxidants which action is based on hydrogen transference reactions [28], by ABTS method, any compound which may be even not real antioxidant like sugars and citric acid but having redox potential lower than ABTS radical-cation can react [29] Depending on the obtained results, it is very difficult to make an unequivocal and undisputable conclusion which of the extraction solvent is the best to obtain as many natural antioxidants from *E. globulus* leaves as possible and to provide an extract rich in antioxidants for manufacture of dietary supplements and medicinal products.

3.3. Neuroprotective activity

Since SH-SY5Y cells remain properties of innate neurons [30], we used this cell model to evaluate the potential neuroprotective effect of *E. globulus* extracts.

Initially, we investigated whether 24 h of treatment with any concentration from $5 \mu\text{g mL}^{-1}$ to $1000 \mu\text{g mL}^{-1}$ of extracts affected SH-SY5Y cell viability (Fig. 1A). MTT assay showed significantly cytotoxic effects when cells were treated with acetone extract at concentrations above $100 \mu\text{g mL}^{-1}$, methanol extract above $250 \mu\text{g mL}^{-1}$ concentration and ethanol extract above $500 \mu\text{g mL}^{-1}$ concentration. Therefore, non-cytotoxic concentrations [from $5 \mu\text{g mL}^{-1}$ to $50 \mu\text{g mL}^{-1}$ for acetone extract; from $5 \mu\text{g mL}^{-1}$ to $100 \mu\text{g mL}^{-1}$ methanol




Fig. 2 – Effects of *Eucalyptus globulus* extracts on ROS production. Human neuroblastoma SH-SY5Y cells were pretreated with non-cytotoxic concentrations of extracts for 24 h, prior to the toxic agent hydrogen peroxide (0.1 mM, 30 min). The intracellular ROS generation was measured by 2',7'-dichlorofluorescein diacetate (DCFH-DA) assay. Each point represents mean values \pm SEM of three independent experiments. * $p < 0.05$ compared with control; # $p < 0.05$ compared with hydrogen peroxide.

extract, and from $5 \mu\text{g mL}^{-1}$ to $250 \mu\text{g mL}^{-1}$ for ethanol extract] were chosen to study the potential protective effect of *E. globulus* extracts against H₂O₂ toxicity.

Hydrogen peroxide is a byproduct of aerobic metabolism involved in many relevant biological and physiological processes, but at high concentrations it also plays a role as inducer of oxidative stress by producing ROS. Consistent studies determined that H₂O₂ is formed during early stages of aggregation of amyloid-beta in AD and of α -synuclein in PD. Substantial accumulation of H₂O₂ and its oxygen-derived free radicals leads to oxidative stress situation [31].

To examine the protective effects of *E. globulus* extracts against H₂O₂ toxicity, SH-SY5Y cells were pretreated with non-cytotoxic concentrations of extracts for 24 h and then H₂O₂ was added at 0.1 mM for 30 min. As shown in Fig. 1B, incubation with H₂O₂ significantly reduced cell viability to 63% compared to control cells. However, pretreatments with extracts led to an increase in cell viability compared to H₂O₂ cells. Particularly, exposure to acetone extract markedly restored cell viability to 77.1% ($5 \mu\text{g mL}^{-1}$), 78.7% ($10 \mu\text{g mL}^{-1}$), 79.8% ($25 \mu\text{g mL}^{-1}$) and 81.6% ($50 \mu\text{g mL}^{-1}$). Moreover, ethanol extract significantly increased cell viability to 92.1% ($5 \mu\text{g mL}^{-1}$), 89.1% ($10 \mu\text{g mL}^{-1}$), 88.5% ($25 \mu\text{g mL}^{-1}$), 85.8% ($50 \mu\text{g mL}^{-1}$) and 79.7% ($100 \mu\text{g mL}^{-1}$). Furthermore, pretreatments with methanol extract exerted significant protective effect at $5 \mu\text{g mL}^{-1}$, $10 \mu\text{g mL}^{-1}$ and $25 \mu\text{g mL}^{-1}$ concentrations by increasing cell viability to 85.1%, 88.9% and 81.5%, respectively. Therefore for




Fig. 3 – Effects of *Eucalyptus globulus* extracts on oxidative stress markers. Human neuroblastoma SH-SY5Y cells were pretreated with non-cytotoxic concentrations of extracts for 24 h, prior to the toxic agent hydrogen peroxide (0.1 mM, 30 min). (A) Effects on glutathione levels, as expressed by ratio GSH/GSSG ratio. (B) Effects on lipid peroxidation, as expressed by thiobarbituric acid reactive substances percentage (TBARS). Each point represents mean values \pm SEM of three independent experiments. * $p < 0.05$ compared with control; # $p < 0.05$ compared with hydrogen peroxide; ^a $p < 0.05$ compare with acetone 25 $\mu\text{g mL}^{-1}$; ^b $p < 0.05$ compare with acetone 50 $\mu\text{g mL}^{-1}$; ^c $p < 0.05$ compare with ethanol 5 $\mu\text{g mL}^{-1}$; ^d $p < 0.05$ compare with ethanol 10 $\mu\text{g mL}^{-1}$; ^e $p < 0.05$ compare with methanol 5 $\mu\text{g mL}^{-1}$.

further studies, we have selected the concentrations 25 and 50 $\mu\text{g mL}^{-1}$ for acetone extract and, 5 and 10 $\mu\text{g mL}^{-1}$ for ethanol and methanol extracts.

Since intracellular ROS overproduction contributes to cell damage and lead to oxidative stress situation, we investigated whether *E. globulus* extracts could protect from H₂O₂-induced ROS generation. Compared with control cells, H₂O₂ generated significant ROS increase (145%). However, all concentrations of extracts markedly decreased ROS generation in H₂O₂-treated SH-SY5Y cells. Particularly, acetone extracts (25 and 50 $\mu\text{g mL}^{-1}$) were the most active, balancing intracellular ROS production by scavenging action (Fig. 2). These results support previous works which revealed that *E. globulus* leaves water extract possess significant ability to ROS and RNS scavenge [32].

Following, we studied the effect of *E. globulus* extracts on cellular redox status markers GSH/GSSG ratio and TBARS level. GSH is the most effective and abundant antioxidant defense mechanism in human body. Previous studies have shown that patients with AD and PD present decreased GSH levels [33]. As shown in Fig. 3A, when SH-SY5Y cells were treated with H₂O₂, GSH/GSSG ratio was declined compared to control cells (GSH/GSSG = 4.1 control cells versus GSH/GSSG = 1.80 H₂O₂-treated neurons) due to low GSH levels and high GSSG levels. However, pretreatments with extracts reversed such drop in GSH/GSSG ratio. Particularly, it highlights methanol extracts (GSH/GSSG = 4.8 at 5 $\mu\text{g mL}^{-1}$ and GSH/GSSG = 6.3 at 10 $\mu\text{g mL}^{-1}$) and ethanol extract at 5 $\mu\text{g mL}^{-1}$ (GSH/GSSG = 4.9).

Lipid peroxidation is an early event in oxidative stress-related neurodegenerative diseases [34]. Then, we investigated the effect of extracts on lipid peroxidation by measuring TBARS content. As shown in Fig. 3B, H₂O₂ treatment significantly increased TBARS levels up to 391.9% fold.

When SH-SY5Y cells were pretreated with extracts, lipid peroxidation was significantly decreased compared to H₂O₂-treated cells. The maximum protective effect against lipid peroxidation was observed for ethanol extract at 5 $\mu\text{g mL}^{-1}$ concentration (151.4%). These findings are in accordance with previous works that demonstrated that *E. globulus* notably decreased plasma and liver malondialdehyde, another product of lipid peroxidation, in streptozotocin-induced diabetic rats [35].

CAT, SOD, GR and GPx antioxidant enzymes play a key role in ROS detoxification and therefore, in prevention and protection against the resulting oxidative stress damage. Previous studies have reported that the neuroprotective mechanism of action of plant phenolic extracts could result from the upregulation of these antioxidant enzymes. As shown in Fig. 4A–D, H₂O₂ significantly reduced CAT, SOD, GPx and GR enzymes activity in SH-SY5Y cells by 59.9%, 33.5%, 23.2% and 26.3%, respectively. On the contrary, pretreatments with extracts increased antioxidants enzymes activities compared to H₂O₂-treated cells. The highest protective effect was observed for all extracts at their higher concentration assayed. Previous works have shown that the aqueous-ethanol leaves extract of *E. globulus* avoid the loss of CAT, SOD and GPx activities in the kidney of rats treated with acetaminophen [36].

4. Conclusions

E. globulus leaves extracts are an important source of antioxidant compounds. The antioxidant potency of acetone, ethanol and methanol extracts depends on the *in vitro* techniques. Moreover, UPLC-ESI-MS/MS analysis profile revealed that these three extracts are rich in phenolic compounds




Fig. 4 – Effects of *Eucalyptus globulus* extracts on antioxidant enzymes. Human neuroblastoma SH-SY5Y cells were pretreated with non-cytotoxic concentrations of extracts for 24 h, prior to the toxic agent hydrogen peroxide (0.1 mM, 30 min). (A) Effects on catalase (CAT) activity; (B) Effects on superoxide dismutase (SOD) activity; (C) Effects on glutathione reductase (GR) activity and (D) Effects on glutathione peroxidase (GPx) activity. Each point represents mean values \pm SEM of three independent experiments. * $p < 0.05$ compared with control; # $p < 0.05$ compared with hydrogen peroxide; ^a $p < 0.05$ compare with acetone 25 $\mu\text{g mL}^{-1}$; ^b $p < 0.05$ compare with acetone 50 $\mu\text{g mL}^{-1}$; ^c $p < 0.05$ compare with ethanol 5 $\mu\text{g mL}^{-1}$; ^d $p < 0.05$ compare with ethanol 10 $\mu\text{g mL}^{-1}$; ^e $p < 0.05$ compare with methanol 5 $\mu\text{g mL}^{-1}$.

(methanol > acetone > ethanol), being chlorogenic acid the major compound identified. Furthermore, *E. globulus* extracts have been demonstrated to prevent from H₂O₂-induced oxidative stress damage by increasing cell viability, GSH levels and antioxidant enzymes activity and, by decreasing ROS production and lipid peroxidation in SH-SY5Y cells. Therefore, these extracts are a potential source of antioxidant compounds with phenolic nature that possess health benefits against oxidative stress, being able to use as raw material for nutritional, food and pharmaceutical supplements.

Conflicts of interest

Authors declare no conflict of interest.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.jfda.2018.05.010>.

REFERENCES

- [1] Barbosa LC, Filomeno CA, Teixeira RR. Chemical variability and biological activities of *Eucalyptus* spp. essential oils. *Molecules* 2016;21:E1671.
- [2] European Medicines Agency. Assessment report on *Eucalyptus globulus* Labill, *Eucalyptus polybractea* R.T. Baker and/or *Eucalyptus smithii* R.T. Baker, aetheroleum EMA/HMPC/307782/2012.
- [3] Gadoth N, Göbel HH. Oxidative stress and free radical damage in neurology. New York: Springer Science+Business Media LLC; 2010.
- [4] Losada-Barreiro S, Bravo-Díaz C. Free radicals and polyphenols: the redox chemistry of neurodegenerative diseases. *Eur J Med Chem* 2017;133:379–402.
- [5] FDA. Draft guidance for industry: bioanalytical method validation (revised). [Online]. US Department of Health and Human Services; September 2013. <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM368107.pdf>.
- [6] Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *JFST* 1995;28:25–30.

- [7] Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999;26:1231–7.
- [8] Asghar MN, Khan IU. Measurement of antioxidant activity with trifluoperazine dihydrochloride radical cation. *Braz J Med Biol Res* 2008;41:455–61.
- [9] Benzie IFF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of antioxidant power: the FRAP assay. *Anal Chem* 1996;39:70–6.
- [10] Apak R, Güçlü K, Demirata B, Özyürek M, Çelik SE, Bektaşoğlu B, et al. Comparative evaluation of various total antioxidant capacity assays applied to phenolic compounds with the CUPRAC assay. *Molecules* 2007;12:1496–547.
- [11] Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Meth* 1983;65:55–63.
- [12] LeBel CP, Ischiropoulos H, Bondy SC. Evaluation of the probe 2',7'-dichlorofluorescein as an indicator of reactive oxygen species formation and oxidative stress. *Chem Res Toxicol* 1992;5:227–31.
- [13] Hissin PJ, Hilf RA. Fluorometric method for determination of oxidized and reduced glutathione in tissues. *Anal Biochem* 1976;74:214–26.
- [14] Mihara M, Uchiyama M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Anal Biochem* 1978;86:271–8.
- [15] Aebi H. Catalase in vitro. *Meth Enzymol* 1984;105:121–6.
- [16] Neha J, Mishra R. Antioxidant activity of Trikatu mega. *Int J Pharm Biomed Res* 2011;2:624–8.
- [17] Staal GE, Helleman PW, de Wael J, Veeger C. Purification and properties of an abnormal glutathione reductase from human erythrocytes. *Biochim Biophys Acta* 1969;185:63–9.
- [18] Wilson SR, Zucker PA, Ruey Ruy CH, Spector A. Development of synthetic compounds with glutathione peroxidase activity. *J Am Chem Soc* 1989;111:5936–9.
- [19] Do QD, Angkawijaya AE, Tran-Nguyen PL, Huynh LH, Soetaredjo FE, Ismadji S, et al. Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *JFDA* 2014;22:296–302.
- [20] Kawabata K, Mukai R, Ishisaka A. Quercetin and related polyphenols: new insights and implications for their bioactivity and bioavailability. *Food Function* 2015;6:1399–417.
- [21] Panche AN, Diwan AD, Chandra SR. Flavonoids: an overview. *JNS* 2016;5:e47.
- [22] Upadhyay R, Mohan Rao LJ. An outlook on chlorogenic acids-occurrence, chemistry, technology, and biological activities. *Crit Rev Food Sci Nutr* 2013;53:968–84.
- [23] Ozcan T, Akpınar-Bayazit A, Yılmaz-Ersan L, Delikanlı B. Phenolics in human health. *IJCEA* 2014;5:393.
- [24] Prior RL. Fruits and vegetables in the prevention of cellular oxidative damage. *Am J Clin Nutr* 2003;78:570S–8S.
- [25] Prior RL, Wu X, Schaich K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J Agr Food Chem* 2005;53:4290–302.
- [26] Lemańska K, Szymusiak H, Tyrakowska B, Zieliński R, Soffers AE, Rietjens IM. The influence of pH on antioxidant properties and the mechanism of antioxidant action of hydroxyflavones. *Free Radic Biol Med* 2001;31:869–81.
- [27] Ozgen M, Reese RN, Tulio AZ, Scheerens JC, Miller AR. Modified 2, 2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) method to measure antioxidant capacity of selected small fruits and comparison to ferric reducing antioxidant power (FRAP) and 2, 2'-diphenyl-1-picrylhydrazyl (DPPH) methods. *J Agr Food Chem* 2006;54:1151–7.
- [28] Staško A, Brezová V, Biskupič S, Mišík V. The potential pitfalls of using 1, 1-diphenyl-2-picrylhydrazyl to characterize antioxidants in mixed water solvents. *Free Rad Res* 2007;41:379–90.
- [29] Pulido R, Bravo L, Saura-Calixto F. Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay. *J Agr Food Chem* 2000;48:3396–402.
- [30] Xie HR, Hu LS, Li GY. SH-SY5Y human neuroblastoma cell line: in vitro cell model of dopaminergic neurons in Parkinson's disease. *Chin Med J* 2010;123:1086–92.
- [31] Chen X, Guo C, Kong J. Oxidative stress in neurodegenerative diseases. *NRR* 2012;7:376–85.
- [32] Almeida IF, Fernandes E, Lima JL, Valentão P, Andrade PB, Seabra RM, et al. Oxygen and nitrogen reactive species are effectively scavenged by *Eucalyptus globulus* leaf water extract. *J Med Food* 2009;12:175–83.
- [33] Gu F, Chauhan V, Chauhan A. Glutathione redox imbalance in brain disorders. *Curr Opin Clin Nutr Metab Care* 2015;18:89–95.
- [34] Bradley-Whitman MA, Lovell MA. Biomarkers of lipid peroxidation in Alzheimer disease: an update. *Arch Toxicol* 2015;89:1035–44.
- [35] Nakhaee A, Bokaeian M, Saravani M, Farhangi A, Akbarzadeh A. Attenuation of oxidative stress in streptozotocin-induced diabetic rats by *Eucalyptus globulus*. *Indian J Clin Biochem* 2009;24:419–25.
- [36] Dhibi S, Mbarki S, Elfeki A, Hfaiedh N. *Eucalyptus globulus* extract protects upon acetaminophen-induced kidney damages in male rat. *BJBMS* 2014;14:99–104.