

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.jfda-online.com

Original Article

Antcins, triterpenoids from *Antrodia cinnamomea*, as new agonists for peroxisome proliferator-activated receptor α



Yu-Jen Wang^a, Shih-Chin Lee^a, Chun-Hua Hsu^b, Yueh-Hsiung Kuo^{c,d,e}, Chien-Chih Yang^a, Fu-Jung Lin^{a,f,*}

^a Department of Biochemical Science and Technology, National Taiwan University, Taipei, Taiwan

^b Department of Agricultural Chemistry, National Taiwan University, Taipei, Taiwan

^c Department of Chinese Pharmaceutical Sciences and Chinese Medicine Resources, China Medical University, Taichung, Taiwan

^d Department of Biotechnology, Asia University, Taichung, Taiwan

^e Chinese Medicine Research Center, China Medical University, Taichung, Taiwan

^f Research Center for Development Biology and Regenerative Medicine, National Taiwan University, Taipei, Taiwan

ARTICLE INFO

Article history:

Received 28 September 2018

Received in revised form

8 November 2018

Accepted 21 November 2018

Available online 4 December 2018

Keywords:

Antrodia cinnamomea

Drug discovery

Metabolism

Nuclear receptor

Peroxisome proliferator-activated receptor α

ABSTRACT

Peroxisome proliferator-activated receptor α (PPAR α) is a nuclear hormone receptor that transcriptionally regulates lipid metabolism and inflammation; therefore, PPAR α full agonists are promising agents to treat dyslipidemia and metabolic disorders. PPAR α full agonists, such as fibrates, are effective anti-hypertriglyceride agents, but their use is limited by adverse side effects. Hence, the aim of this study was to identify small molecules that can activate PPAR α while minimizing the adverse effects. *Antrodia cinnamomea*, a rare medical mushroom, has been used widely in Asian countries for the treatment of various diseases, including liver diseases. Antcin B, H and K (antcins) and ergostatrien-3 β -ol (EK100) are bioactive compounds isolated from *A. cinnamomea* with anti-inflammatory actions. Antcins, ergostane-type triterpenoids, contain the polar head with carboxylate group and the sterol-based body. Here, we showed at the first time that sterol-based compounds, antcins, but not EK100, activate PPAR α in a cell-based transactivation study. The *in silico* docking studies presented several significant molecular interactions of antcins, including Tyr314, and His440 in the ligand-binding domain of PPAR α , and these interactions are required for helix 12 (H12) stabilization. We propose that PPAR α activation activity of antcins is related to their binding mode which requires conventional H12 stabilization, and that antcins can be developed as safe selective PPAR α modulators.

Copyright © 2018, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Abbreviations: *A. cinnamomea*, *Antrodia cinnamomea*; ALP, alkaline phosphatase; EK100, ergostatrien-3 β -ol; LBD, Ligand Binding Domain; MetS, metabolic syndrome; PPARs, peroxisome proliferator-activated receptors; TZDs, thiazolidinediones; T2DM, type 2 diabetes mellitus.

* Corresponding author. Department of Biochemical Science and Technology, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd., Taipei, 10617, Taiwan.

E-mail address: fujlin@ntu.edu.tw (F.-J. Lin).

<https://doi.org/10.1016/j.jfda.2018.11.004>

1021-9498/Copyright © 2018, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Peroxisome proliferator-activated receptors (PPARs) are members of the nuclear receptor superfamily of ligand-activated transcription factors. There are three PPAR iso-types, PPAR α (NR1C1, encoded by *PPARA*), PPAR β/δ (NR1C2, encoded by *PPARD*), and PPAR γ (NR1C3, encoded by *PPARG*) [1]. In general, all three members of the PPAR subfamily function as lipid sensors that transcriptionally modulate metabolic programs involved in lipid and energy metabolism. PPAR has been proposed as an important therapeutic target for metabolic complications, such as dyslipidemia, diabetes and fatty liver disease.

Specifically, each member has its tissue-specific distribution patterns and ligand specificities, which contribute to their distinct biological functions. PPAR α is highly expressed in liver, where it has crucial roles in regulation of adaptive response to fasting by controlling fatty acid transport, β -oxidation and ketogenesis [2]. Mice lacking PPAR α are prone to develop to liver steatosis, inflammation, and hepatocellular carcinoma [3–5]. Various fatty acids and their metabolites are considered as biological ligands for PPAR α . It also can be activated by the fibrate class of anti-lipidemic drugs, which includes fenofibrate and clofibrate [6]. PPAR γ is highly expressed in adipose tissues, where it governs adipogenesis and adipocyte function [7]. PPAR γ is activated by thiazolidinediones (TZDs), such as rosiglitazone and pioglitazone. It has been suggested that TZDs function as insulin-sensitizers and are widely used for the treatment of type 2 diabetes mellitus (T2DM) [8,9].

Antrodia cinnamomea is a rare and precious medical mushroom found in Taiwan and has been reported to have anti-inflammatory, anticancer, hepatoprotective and immunomodulating properties [10]. Thus, it is used widely in Asian countries for the treatment of various diseases, including liver disease. A few bioactive compounds have been identified in fruiting bodies and mycelia of *A. cinnamomea*, including triterpenoids, polysaccharide, succinic and maleic acid derivatives, and ubiquinone derivatives [10,11]. Antcins are the typical triterpenoid compounds of *A. cinnamomea*. Among them, both antcin B and H possess anti-inflammatory, anticancer, insecticidal and hepatoprotective activities [12–17]. Similarly, antcin C and H have anti-inflammatory and anticancer activities *in vitro* [16,18–20]. Ergostatrien-3 β -ol (EK100), a major component of the submerged whole broth of *A. cinnamomea*, was reported to have anti-inflammatory and anti-photodamaging activities in mice [21]. In addition to anti-inflammatory effects, a recent study showed that antcin K and EK100 displayed antihyperlipidemic and antidiabetic effects in a diet-induced obesity mouse model [22,23]. This prompts us to investigate if antcins and EK100 mediate lipid and glucose-lowering effects presumably through activation of PPARs.

In the present study, we examined the effects of antcin B, H, K and EK100 on PPAR α and PPAR γ activation using a cell-based transactivation assay, and predicted the potential nuclear receptor–ligands interactions using *in silico* docking approaches. As modest PPAR α agonists, antcin B, H, and K selectively activated PPAR α activity. Computational studies

revealed that antcins occupied the ligand-binding site of PPAR α like PPAR α agonist Wy-14,643, which requires AF2 helix stabilization. Together, our findings suggest that natural compounds antcins are novel PPAR α activators, and would be potential chemicals for the improvement of current anti-dyslipidemic agents.

2. Methods

2.1. Chemicals and reagents

The freeze-dried mycelium of *A. cinnamomea* of the submerged whole broth was extracted with methanol. Antcin B, H, and K were isolated from the fruiting body of *A. cinnamomea*. The detailed purification process of EK100, antcin B, H, and K were described previously [24,25].

2.2. Cell culture and treatment

Chinese hamster ovary cells (CHO–K1) (ATCC CRL-9618) were maintained in Ham's F12 Nutrient medium (Gibco) supplemented with 10% heat-inactivated fetal bovine serum (FBS) (Gibco) at 37 °C in a humidified environment containing 5% CO₂. Cells were subcultured every 3 days in 25- or 75-cm² flasks. All chemicals were dissolved in absolute ethanol (Sigma, St. Louis, MO) and were then added at various concentrations to cells for 48 h. The final ethanol concentration was less than 0.1%.

2.3. Expression and reporter vectors

The chimeric receptor constructs used were pBK-CMV Gal4-rPPAR α ligand binding domain (Gal4-rPPAR α LBD) and the pBK-CMV Gal4-rPPAR γ ligand binding domain (Gal4-rPPAR γ LBD), respectively. The reporter gene was pBK-CMV-(UAS)₄-tk-alkaline phosphatase (ALP) [26].

2.4. Transient transfection and alkaline phosphatase (ALP) activity assay

CHO–K1 cells were seeded in a 96-well plate (2.5 $\times$ 10⁴ cells per well) in Ham's F12 Nutrient medium supplemented with 10% charcoal-stripped FBS and incubated overnight before transfection. Then, nearly confluent cells were transfected with 0.1 μ g the chimeric receptor constructs containing PPAR LBD and 0.04 μ g of the reporter plasmid using Lipofectamine 2000 (Thermo Fisher Scientific, Waltham, MA) according to the manufacturer's instructions. Five hours after transfection, the supernatants were removed. The transfected cells were then treated for 48 h with Wy-14,643 (Cayman Chemicals, Ann Arbor, MI), BRL 49653 (rosiglitazone) (Sigma, St. Louis, MO), and other reagents at the indicated concentrations. All tested compounds were prepared in Ham's F12 Nutrient medium supplemented with 10% TCM (Protide Pharmaceuticals, Inc, Lake Zurich, IL). The supernatants were collected, and ALP activity was measured by adding SEAP buffer, and p-Nitrophenyl Phosphate (pNPP) (Sigma, St. Louis, MO) was used as the substrate. Absorbance at 405 nm was measured using a Multiskan GO Microplate Reader (Thermo Fisher Scientific,

Waltham, MA). Relative PPAR α or PPAR γ activity is expressed as a fold change relative to the vehicle control.

2.5. MTT assay

To determine the effect of the chemicals on cell viability, cells were applied with 55 μ L of 0.45 mg/mL 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma, St. Louis, MO) dissolved in Ham's F12 Nutrient medium in the dark, and then cells were incubated for 3 h at 37 °C, 5% CO₂. To solubilize the product of MTT cleavage, 100 μ L of isopropanol containing 0.04 N HCl was added to each well and thoroughly mixed using a microplate shaker (Scientific Industries, Inc., Bohemia, NY). After 20 min of HCl-isopropanol addition, absorbance at 540 nm was measured using a Multiskan GO Microplate Reader. Relative cell viability is expressed as percentage change relative to the vehicle control.

2.6. Docking simulation

Crystal structures of the ligand-bound LBD of human PPAR α (PDB ID: 4BCR) in complex with Wy-14,643 were employed as receptor for docking study, which was edited using PyMOL (Schrödinger, NY) to select a single polypeptide chain and remove all water molecules and co-crystallized ligands. To perform docking with AutoDock4 [27], the structures of receptor and ligands were transformed to the pdbqt file format by using AutoDock Tools package. A grid with dimensions of 25 $\times$ 25 $\times$ 25 points was centered to ensure coverage of the binding pocket of LBD. The structures containing selected pose of ligands upon LBD were applied a procedure of solvation and following energy minimization to obtain the final structures. 2-Dimensional (2D) diagrams of interacting residues were depicted by default procedures and tools in Discovery Studio 3.5 software (Accelrys Inc., San Diego, CA).

2.7. Statistical analysis

Data are presented as the means $\pm$ SD. Each experiment was performed in triplicate. Statistical analyses were carried out

using IBM SPSS Statistics 20 (IBM Corporation, USA). One-way analysis of variance (ANOVA) was used to determine differences among control and experimental groups. $P < 0.05$ was considered statistical significant. EC₅₀ values were calculated with GraphPad Prism (GraphPad Software, San Diego, CA).

3. Results

3.1. Activation of PPAR α ligand binding domains (LBD) by Wy-14,643 and antcins in CHO–K1 cells

Antcin B, H, K and EK100 contain steroid-like four rings structure, and a side chain. All antcins and Wy-14,643, but not EK100 contain a carboxylic acid moiety on the side chain (Fig. 1). As all test compounds were dissolved in ethanol, we first examined the cytotoxic effects of the organic solvent on CHO–K1 cell by MTT assay. There were little or no effects on cell viability when the cells were treated with 1% (v/v) of ethanol, or lower than this concentration (Fig. S1) (Supplementary material). 10% ethanol significantly decreased cell survival to 15.7%. Next, we measured the effects of test compounds at various concentrations (0.1–200 μ M) on the viability of CHO–K1 cells using MTT assay. None of the compounds caused a reduction in cell viability to 20% of the control when applied up to a level of 10 μ M, except for 10 μ M of EK100, which caused a 45% reduction in cell viability (Fig. 2A).

To understand whether the test compounds activate PPAR α , PPAR α activities were determined by the ALP reporter assay using co-transfection of a plasmid carrying chimera genes encoding the Gal4 PPAR α ligand-binding domain (LBD), and a plasmid carrying UAS ALP reporter gene as described previously [28]. Cells were treated with ethanol (vehicle), Wy-14,643 (PPAR α agonist), or test compounds for 48 h. The levels of PPAR α activation were expressed as a fold change relative to the vehicle control. As shown in Fig. 2B, the strongest activation was observed following the application of Wy-14,643, which had an EC₅₀ value of 0.08 μ M. In this system, antcin B, H and K activated PPAR α activity. As estimated from concentration–response curves, the EC₅₀ for antcin H was




Fig. 1 – Chemical structure of (A) Wy-14,643, (B) EK100, (C) antcin B, (D) antcin H, and (E) antcin K.




Fig. 2 – Antcins are PPAR α agonist. (A) Cell viability and (B) PPAR α transactivation by different concentrations of Wy-14,643, antcin B, H, and K, and EK100. The CHO–K1 cells were co-transfected with the plasmids encoding chimeric receptor constructs containing PPAR α LBD and their cognate ALP reporters (see “Methods”). After transfection, cells were treated with vehicle (ethanol) or ligands with various concentrations. Data are means $\pm$ SD of at least three replicates.

found to be 9.3 μ M. The EC₅₀ values for antcin B and K in this system could not be calculated since activation could not reach top plateau at the range of tested concentrations of these substances.

All of the examined antcins activated PPAR α in a dose-dependent manner (Fig. 2B). Expression of the results of the application of 10 μ M of each compound as a percentage of the control revealed a trend, in which the order of activation capacity was antcin B = antcin H > antcin K, with calculated folds of 2.5, 2.5 and 1.1, respectively (Fig. 2B). As concentrations were increased, maximal activation (5.7-fold) was achieved by application of 200 μ M of antcin K. This activation was better than that observed for Wy-14,643, a well-recognized specific activator of PPAR α (Fig. 2B).

3.2. Little or no activation of PPAR γ LBD by antcin B and H in CHO–K1 cells

To understand whether antcin B and H activate PPAR γ in CHO–K1 cells, cells were transfected with the PPAR γ LBD plasmid construct and reporter construct and treated with BRL 49,653 (PPAR γ agonist) for 48 h. 1 μ M BRL49653 activated PPAR γ more than 7 folds compared with vehicle in CHO–K1 cells, whereas none of antcins activated PPAR γ activity at concentrations up to 30 μ M (Fig. 3A). Cell viability was

unaffected upon antcins treatment (Fig. 3B). These results suggest that antcins are less likely to act as PPAR γ agonists.

3.3. Computational analyses of bound ligands

The orientation and residue bonding of Wy-14,643 inferred from the X-ray structure of human PPAR α (PDB code 4BCR) served as a model for computing the interactions of Wy-14,643 and antcins. The overall structure of LBD of human PPAR α (here denoted hPPAR α _LBD) encompasses residues 196–467, which consists of 12 helices and a 3-stranded antiparallel β -sheet (Fig. 4A). C-terminal helix (helix 12 or AF-2 helix) which is packed onto the core of LBD is crucial for maintaining the active conformation of PPAR α [29]. The region between helix 3 and the 3-stranded antiparallel β -sheet is proposed to be ligand entry site [29]. Based on the structural analysis, there are three main features can be described at the binding site, including hydrogen bonds, a hydrophobic region, and electrostatic interactions. Residues Ser280 (H3), Tyr314 (H5), and Tyr464 (H12) form a hydrogen bond network with the carboxylic acid group of the Wy-14,643 molecule (Figs. 4C and 5A). In addition, Wy-14,643 also makes hydrophobic contacts with residues Phe318, Leu321, Met330, Leu331, and Leu460 (Fig. 5A).

Next, we compared the *in silico* docking results of antcins and Wy-14,643. Intriguingly, antcins interact with the hPPAR α _LBD like Wy-14,643 did (Fig. 4B, D, E and F). The carboxylic acid moiety of antcin H forms a hydrogen bond with residues Tyr314 on helix 5 (Fig. 4D), while the carboxylic acid moiety of antcin B and K form hydrogen bonds with residues His440 on helix 10 of hPPAR α _LBD (Fig. 4E and F). As identified for other PPAR α agonist, the hydrogen bonds linking agonist and Ser280, Tyr314, His440 and Tyr464 of the receptor are crucial for stabilizing AF2 helix in the active conformational state [29].

The antcins-nuclear receptor interaction involved additional interactions at the binding site. Since antcins are composed of steroid-like four rings and are much bulkier than Wy-14,643, it is capable of carrying out a significant part of the hydrophobic interactions, such as those between antcins and residues Asn219, Met220, Phe273, Val324, and Val444 (Fig. 5A). Due to the structure similarity, antcin B, H, and K shared interactions with 20 surrounding residues of hPPAR α _LBD (Fig. 5A and B). Among them, antcins and Wy-14,643 commonly interacted with 10 PPAR α residues, including Cys276, Gln277, Thr279, Ser280, Tyr314, Met330, Met355, His440, Leu460, and Tyr464 (Fig. 5B, highlighted in bold in the box).

Combining the hydrophobicity analysis plot of PPAR α ligand binding pocket and the common residues shared by antcins and Wy-14,643, we then divided these 10 residues into two groups. One group includes the seven residues Gln277, Thr279, Ser280, Tyr314, His440, Leu460 and Tyr464, which contributes to the hydrophilic center near the H12 of the pocket (Fig. 6A). The other group contains the rest residues Cys276, Met330 and Met355, which comprise the hydrophobic surface between H7 and β sheets (Fig. 6B).

Among the tested antcins, antcin H had the lowest EC₅₀ value. We further examined the structure-activity relationships of antcins. Antcin B has a carbonyl group on A ring, while antcin H and K contain one and two hydroxyl groups on




Fig. 3 – Antcin B and H are not PPAR γ agonists. (A) PPAR γ transactivation and (B) Cell viability by different concentrations of antcin B and H. The CHO–K1 cells were co-transfected with the plasmids encoding chimeric receptor constructs containing PPAR γ LBD and their cognate ALP reporters (see “Methods”). After transfection, cells were treated with vehicle (ethanol) or ligands with various concentrations. Data are means $\pm$ SD of at least three replicates.

A ring, respectively (Fig. 1). We found that among the residues of ligand-binding pocket surrounding A ring, Asn219, Glu282, and Glu286 contribute to a local polar center, whereas Met220, Val324, and Leu331 make up the other non-polar region (Fig. 6C). Antcin K has two hydroxyl groups on ring A. One of them was pointed to the direction of the polar center, while the other was headed toward the non-polar hemisphere

(Fig. 6C). The possible repulsion may lead to the decreased potency of antcin K. In contrast, the hydroxyl group on A ring of antcin H was predicted to pointed to the direction of the polar center, it may provide the stabilization energy for binding of antcin H to the ligand-binding site (Fig. 6C). This may account for the lowest EC₅₀ of antcin H among antcins. In addition to interactions around ring A, the hydroxyl or carbonyl groups on the others rings of antcins can engage in additional hydrophobic interactions at the binding site. For instance, antcins were predicted to make hydrophobic interactions with Leu321 through their carbonyl group or hydroxyl group on B ring (Fig. 5A).

4. Discussion

Increasing evidence has shown that PPAR α plays a crucial role in fatty acid metabolism, as it controls the expression of genes mediated peroxisomal and mitochondrial β -oxidation, ω -hydroxylation, and ketogenesis [30]. PPAR α has been considered as an important therapeutic target for metabolic disorders, such as dyslipidemia, and fatty liver diseases. Hence, identification of PPAR α agonists has a major medical interest. This is the first study showing that natural triterpenoid compounds antcins activate PPAR α .

One of the common structural features among the activators is a carboxylate group, which is not found in the non-activator, EK100. These findings highlight the importance of the carboxylate moieties of these ligands in the activation of PPAR α . Indeed, most known PPAR agonists share features including a carboxylate head group, a central hydrophobic part, and a flexible linker to the tail. The results of the computational analysis suggest that antcins are inserted in a similar orientation as Wy-14,643, in which the carboxylic group faces the hydrophilic region of the binding site and is close to a key switch helix, H12 (AF2 helix) [31]. In the absence of ligand, H12 is highly dynamic and is mobile along with other portions of the LBD [32]. Ligands stabilize the receptor folding and lower the conformational dynamics [32]. Like Wy-14,643, antcins form hydrogen bonds with Tyr314 on helix 5 and His440 on the helix 10 by its carboxylic group. Several studies showed that hydrogen bond network between the carboxyl group of ligands and the residues surrounding AF2 helix can stabilize the dynamics of AF2 helix, which permits coactivator recruitment [33,34]. These specific interactions enhance the transactivation activity of the PPARs [34]. As expected, antcins exhibit PPAR α activating ability in the cell-based transactivation assay. These data suggest that formation of hydrogen bonds between residues near H12 of PPAR α and its acidic ligands is crucial for ligand activation ability.

A recent study reported a bipartite mechanism of PPAR α activation by Wy-14,643, which involves two ligands in the pocket simultaneously [31]. The first Wy-14,643 occupies standard PPAR α agonist position, and requires conventional direct H12 stabilization [31]. The second Wy-14,643 molecule binds to a site positioned between H2' and H3 (hereafter denoted site two), and is primarily stabilized by nonpolar interactions [31]. Site two is located in the region called omega loop. Activity assay showed that site two is required for full Wy-14,643 activity [31]. According to our results of docking




Fig. 4 – Three-dimensional structures of PPAR α -LBD stabilized by the ligands. (A, B) Overall structure of Wy-14,643 (A) and antcin H (B) bound with PPAR α in ribbon representation. The bound ligands are shown in stick representation. The H12 (AF2) helix is labeled. Missing residues in the H2–H2' loop are depicted as an orange curve. (C–F) Close-up view of the Wy-14,643 (C), antcin H (D), antcin B (E), and antcin K (F) at the binding site of PPAR α showing the interacting residues. Ligands are shown as black sticks, and receptor residues are shown as grey sticks. The bound ligands and receptor residues are shown in stick representation with oxygen, nitrogen, chloride, sulfur and hydrogen atoms depicted in red, blue, green, yellow, and white, respectively. Hydrogen bonds are shown as dotted green lines.

simulation, all antcins and EK100 can be docked to the standard PPAR α agonist position (Fig. 4 and data not shown), but not all of them induce PPAR α activation *in vitro*. One possibility is that EK100 is lack of carboxylate group, and the other is that the conventional docked position is not sufficient to induce PPAR α activation. As antcins, which contain a carboxylate head group, have higher EC₅₀ values than Wy-14,643, it is tempting to speculate that site two and/or simultaneous insertion of two ligands into the pocket might be also essential for full activity of the ligands. Further studies with site

mutation analysis need to be performed to decipher this possibility.

In the present study, we showed that antcins activated PPAR α not PPAR γ . PPAR α and PPAR γ share 60–70% sequence identity in their LBDs. Although both subtypes bind to naturally occurring polyunsaturated fatty acids, they exhibit ligand selectivity. For instance, only PPAR α binds to a wide range of saturated fatty acids [35]. This is supported by structural analyses revealing that PPAR α ligand binding pocket has the more hydrophobic nature compared to that of




Fig. 5 – Major interactions established by key residues were identified using docking analysis. (A) Two-dimensional interaction diagram of the predicted binding mode of the ligands (Wy-14,643, antcin B, H, and K), PPAR α , and bound water molecules (blue). Ligands are shown as black sticks with oxygen, nitrogen, chloride, sulfur and hydrogen atoms depicted in red, blue, green, yellow, and white, respectively. Hydrophobic and electrostatic interactions are indicated in green and pink, respectively. Hydrogen bonds are shown as dashed blue lines. (B) Venn diagram showing the overlapping numbers of key residues by antcin B, H and K. Twenty residues shared by antcins were listed in the box, and ten of them commonly interacted by Wy-14,643 were highlighted in bold.

PPAR γ [34]. As expected, antcins containing the steroid core ring structure were detected to make hydrophobic interactions with PPAR α LBD in the software simulation. Furthermore, one of the determinants of selectivity between PPAR α and PPAR γ is the substitution of Ala454 on the hinge

(loop) of H12 in PPAR α for Met463 in PPAR γ [36]. H12 dynamic stabilization is crucial for activation of PPARs by ligands [37]. The larger volume of the Met463 side chain in PPAR γ may act as a lid to limit the accessibility of ligands to ligand binding pocket. Thus, not only hydrophobic nature of PPAR α ligand




Fig. 6 – Maps of the hydrophobicity at the binding site of PPAR α docked with the antcin H and K. (A, B) Ten residues shared by antcins and Wy-14,643 are shown as grey sticks. The bound ligands are shown as black sticks. H3, H5 and H7 of PPAR α are shown in ribbon representation. (C) Maps of the hydrophobicity at the binding site of PPAR α docked with the antcin H and K. Ligands and receptor residues surrounding ring A of antcins are shown as black and yellow sticks, respectively. Regions surrounding ring A are highlighted in dotted rectangles.

binding pocket, but the amino acids on the hinge (loop) of H12 may be involved in the ligand selectivity.

Several studies have demonstrated that PPAR α activation ameliorates metabolic disorder and non-alcoholic fatty liver disease (NAFLD) in diet-induced obesity [1]. In this study, we showed that triterpenoid compounds antcins activate PPAR α *in vitro* and *in silico*. Recent studies showed that the crude extract of *A. cinnamomea*, or purified antcin K prevent obesity and display antihyperlipidemic effects through regulating AMPK signaling in high fat diet-induced obese mice [22,38]. One could speculate that the protective effects of antcins against obesity and hepatic steatosis could be also attributed to their abilities to activate PPAR α . In addition to synthetic agonists, several natural compounds, including triterpenoids, have been identified as PPAR α modulators [39]. For example, the pentacyclic triterpene oleanolic acid was found to stimulate PPAR α activation by a reporter gene

assay using PPAR response element activity in CV-1 keratinocytes cells [40]. Cucurbitane-type triterpenoids isolated from wild bitter melon (*Momordica charantia* L.) were shown to activate PPAR α and PPAR γ in a transactivation assay [26]. In addition, wild bitter gourd has been shown to ameliorate metabolic syndrome (MetS) in animal models and subjects with MetS [26,41]. Thus, natural PPAR α agonists may have therapeutic potential for treating metabolic disorders.

5. Conclusion

In summary, the present study is the first to elucidate molecular aspects of the nuclear receptor activation effects induced by antcins using both *in vitro* and *in silico* approaches. Our findings identified potential PPAR α agonists from herbs, and provided insight into the interaction of antcins with PPAR α pathways.

Conflicts of interest statement

The authors declared that there are no conflicts of interest.

Authors' contributions

Y.-J.W. and S.-C.L. performed the experiments, analyzed and interpreted the data. C.-C.Y. interpreted the data, and provided advice. Y.-H.K. provided the reagents and advice. C.-H.H. and F.-J.L. designed and performed the experiments, interpreted the data, and supervised the study. F.-J.L. wrote the manuscript.

Acknowledgements

We thank Dr. Ching-Jang Huang for helpful discussion. This work was supported by grants MOST 105-2320-B-002-011 (F.-J.L.), MOST 106-2311-B-002-038 (F.-J.L.), MOST 106-2113-M-002-020 (C.-H.H.), MOST 105-2313-B-002-050-MY3 (C.-C.Y.), and MOST 107-2321-B-002-047 (C.-C.Y.) from the Ministry of Science and Technology, Taiwan, and by grant MOHW107-TDU-B-212-123004 (Y.-H.K.) from Taiwan Ministry of Health and Welfare Clinical Trial Center. This work was also supported in part by the “Chinese Medicine Research Center, China Medical University” from the Featured Areas Research Center Program within the framework of the Higher Education Sprout Project by the Ministry of Education (MOE) in Taiwan (CMRC-CHM-4).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jfda.2018.11.004>.

REFERENCES

- [1] Gross B, Pawlak M, Lefebvre P, Staels B. PPARs in obesity-induced T2DM, dyslipidaemia and NAFLD. *Nat Rev Endocrinol* 2017;13:36–49.
- [2] Pawlak M, Lefebvre P, Staels B. Molecular mechanism of PPARalpha action and its impact on lipid metabolism, inflammation and fibrosis in non-alcoholic fatty liver disease. *J Hepatol* 2015;62:720–33.
- [3] Zhang N, Chu ES, Zhang J, Li X, Liang Q, Chen J, et al. Peroxisome proliferator activated receptor alpha inhibits hepatocarcinogenesis through mediating NF-kappaB signaling pathway. *Oncotarget* 2014;5:8330–40.
- [4] Abdelmegeed MA, Yoo SH, Henderson LE, Gonzalez FJ, Woodcroft KJ, Song BJ. PPARalpha expression protects male mice from high fat-induced nonalcoholic fatty liver. *J Nutr* 2011;141:603–10.
- [5] Stienstra R, Mandarm S, Patsouris D, Maass C, Kersten S, Muller M. Peroxisome proliferator-activated receptor alpha protects against obesity-induced hepatic inflammation. *Endocrinology* 2007;148:2753–63.
- [6] Staels B, Maes M, Zamboni A. Fibrates and future PPARalpha agonists in the treatment of cardiovascular disease. *Nat Clin Pract Cardiovasc Med* 2008;5:542–53.
- [7] Willson TM, Lambert MH, Kliewer SA. Peroxisome proliferator-activated receptor gamma and metabolic disease. *Annu Rev Biochem* 2001;70:341–67.
- [8] Soccio RE, Chen ER, Lazar MA. Thiazolidinediones and the promise of insulin sensitization in type 2 diabetes. *Cell Metab* 2014;20:573–91.
- [9] Chana RS, Lewington AJ, Brunskill NJ. Differential effects of peroxisome proliferator activated receptor-gamma (PPAR gamma) ligands in proximal tubular cells: thiazolidinediones are partial PPAR gamma agonists. *Kidney Int* 2004;65:2081–90.
- [10] Geethangili M, Tzeng YM. Review of pharmacological effects of *Antrodia camphorata* and its bioactive compounds. *Evid Based Complement Alternat Med* 2011;2011:212641.
- [11] Zhang BB, Hu PF, Huang J, Hu YD, Chen L, Xu GR. Current advances on the structure, bioactivity, synthesis, and metabolic regulation of novel ubiquinone derivatives in the edible and medicinal mushroom *Antrodia cinnamomea*. *J Agric Food Chem* 2017;65:10395–405.
- [12] Huo Y, Win S, Than TA, Yin S, Ye M, Hu H, et al. Antcin H protects against acute liver injury through disruption of the interaction of c-Jun-N-terminal kinase with mitochondria. *Antioxid Redox Signal* 2017;26:207–20.
- [13] Chiu KY, Chen TH, Wen CL, Lai JM, Cheng CC, Liu HC, et al. Antcin-H isolated from *Antrodia cinnamomea* inhibits renal cancer cell invasion partly through inactivation of FAK-ERK-C/EBP-beta/c-Fos-MMP-7 pathways. *Evid Based Complement Alternat Med* 2017;2017:5052870.
- [14] Yeh CT, Rao YK, Yao CJ, Yeh CF, Li CH, Chuang SE, et al. Cytotoxic triterpenes from *Antrodia camphorata* and their mode of action in HT-29 human colon cancer cells. *Cancer Lett* 2009;285:73–9.
- [15] Male KB, Rao YK, Tzeng YM, Montes J, Kamen A, Luong JH. Probing inhibitory effects of *Antrodia camphorata* isolates using insect cell-based impedance spectroscopy: inhibition vs chemical structure. *Chem Res Toxicol* 2008;21:2127–33.
- [16] Shen YC, Wang YH, Chou YC, Chen CF, Lin LC, Chang TT, et al. Evaluation of the anti-inflammatory activity of zhangkui acids isolated from the fruiting bodies of *Antrodia camphorata*. *Planta Med* 2004;70:310–4.
- [17] Chen CH, Yang SW, Shen YC. New steroid acids from *Antrodia cinnamomea*, a fungal parasite of *Cinnamomum micranthum*. *J Nat Prod* 1995;58:1655–61.
- [18] Tien AJ, Chien CY, Chen YH, Lin LC, Chien CT. Fruiting bodies of *Antrodia cinnamomea* and its active triterpenoid, antcin K, ameliorates N-Nitrosodiethylamine-induced hepatic inflammation, fibrosis and carcinogenesis in rats. *Am J Chin Med* 2017;45:173–98.
- [19] Huang YL, Chu YL, Ho CT, Chung JG, Lai CI, Su YC, et al. Antcin K, an active triterpenoid from the fruiting bodies of basswood-cultivated *Antrodia cinnamomea*, inhibits metastasis via suppression of integrin-mediated adhesion, migration, and invasion in human hepatoma cells. *J Agric Food Chem* 2015;63:4561–9.
- [20] Gokila Vani M, Kumar KJ, Liao JW, Chien SC, Mau JL, Chiang SS, et al. Antcin C from *Antrodia cinnamomea* protects liver cells against free radical-induced oxidative stress and apoptosis in vitro and in vivo through Nrf2-dependent mechanism. *Evid Based Complement Alternat Med* 2013;2013:296082.
- [21] Kuo YH, Lin TY, You YJ, Wen KC, Sung PJ, Chiang HM. Antiinflammatory and antiphotodamaging effects of

- ergostatrien-3 β -ol, isolated from *Antrodia camphorata*, on hairless mouse skin. *Molecules* 2016;21.
- [22] Kuo YH, Lin CH, Shih CC, Yang CS. Antcin K, a triterpenoid compound from *Antrodia camphorata*, displays antidiabetic and antihyperlipidemic effects via glucose transporter 4 and AMP-activated protein kinase phosphorylation in muscles. *Evid Based Complement Alternat Med* 2016;2016:4867092.
- [23] Kuo YH, Lin CH, Shih CC. Ergostatrien-3 β -ol from *Antrodia camphorata* inhibits diabetes and hyperlipidemia in high-fat-diet treated mice via regulation of hepatic related genes, glucose transporter 4, and AMP-activated protein kinase phosphorylation. *J Agric Food Chem* 2015;63:2479–89.
- [24] Shao YY, Chen CC, Wang HY, Chiu HL, Hseu TH, Kuo YH. Chemical constituents of *Antrodia camphorata* submerged whole broth. *Nat Prod Res* 2008;22:1151–7.
- [25] Shen CC, Kuo YC, HR L, Lin LC, Don MJ, Chang TT, et al. New ergostane and lanostane from *Antrodia camphorata*. *J Chin Med* 2003;14:247–58.
- [26] Lu KN, Hsu C, Chang ML, Huang CJ. Wild bitter gourd increased metabolic rate and up-regulated genes related to mitochondria biogenesis and UCP-1 in mice. *J Funct Foods* 2013;5:668–78.
- [27] Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS, et al. AutoDock4 and AutoDockTools4: automated docking with selective receptor flexibility. *J Comput Chem* 2009;30:2785–91.
- [28] Chao CY, Huang CJ. Bitter gourd (*Momordica charantia*) extract activates peroxisome proliferator-activated receptors and upregulates the expression of the acyl CoA oxidase gene in H4IIEC3 hepatoma cells. *J Biomed Sci* 2003;10:782–91.
- [29] Cronet P, Petersen JF, Folmer R, Blomberg N, Sjoblom K, Karlsson U, et al. Structure of the PPAR α and - γ ligand binding domain in complex with AZ 242; ligand selectivity and agonist activation in the PPAR family. *Structure* 2001;9:699–706.
- [30] Schoonjans K, Staels B, Auwerx J. Role of the peroxisome proliferator-activated receptor (PPAR) in mediating the effects of fibrates and fatty acids on gene expression. *J Lipid Res* 1996;37:907–25.
- [31] Bernardes A, Souza PC, Muniz JR, Ricci CG, Ayers SD, Parekh NM, et al. Molecular mechanism of peroxisome proliferator-activated receptor α activation by WY14643: a new mode of ligand recognition and receptor stabilization. *J Mol Biol* 2013;425:2878–93.
- [32] Rastinejad F, Huang P, Chandra V, Khorasanizadeh S. Understanding nuclear receptor form and function using structural biology. *J Mol Endocrinol* 2013;51:T1–21.
- [33] Gampe Jr RT, Montana VG, Lambert MH, Miller AB, Bledsoe RK, Milburn MV, et al. Asymmetry in the PPAR γ /RXR α crystal structure reveals the molecular basis of heterodimerization among nuclear receptors. *Mol Cell* 2000;5:545–55.
- [34] Xu HE, Lambert MH, Montana VG, Plunket KD, Moore LB, Collins JL, et al. Structural determinants of ligand binding selectivity between the peroxisome proliferator-activated receptors. *Proc Natl Acad Sci U S A* 2001;98:13919–24.
- [35] Xu HE, Lambert MH, Montana VG, Parks DJ, Blanchard SG, Brown PJ, et al. Molecular recognition of fatty acids by peroxisome proliferator-activated receptors. *Mol Cell* 1999;3:397–403.
- [36] Carrieri A, Giudici M, Parente M, De Rosas M, Piemontese L, Fracchiolla G, et al. Molecular determinants for nuclear receptors selectivity: chemometric analysis, dockings and site-directed mutagenesis of dual peroxisome proliferator-activated receptors α / γ agonists. *Eur J Med Chem* 2013;63:321–32.
- [37] Rastinejad F, Ollendorff V, Polikarpov I. Nuclear receptor full-length architectures: confronting myth and illusion with high resolution. *Trends Biochem Sci* 2015;40:16–24.
- [38] Peng CH, Yang MY, Yang YS, Yu CC, Wang CJ. *Antrodia cinnamomea* prevents obesity, dyslipidemia, and the derived fatty liver via regulating AMPK and SREBP signaling. *Am J Chin Med* 2017;45:67–83.
- [39] Rigano D, Sirignano C, Tagliatalata-Scafati O. The potential of natural products for targeting PPAR α . *Acta Pharm Sin B* 2017;7:427–38.
- [40] Lee HK, Nam GW, Kim SH, Lee SH. Phytocomponents of triterpenoids, oleanolic acid and ursolic acid, regulated differently the processing of epidermal keratinocytes via PPAR- α pathway. *Exp Dermatol* 2006;15:66–73.
- [41] Tsai CH, Chen EC, Tsay HS, Huang CJ. Wild bitter gourd improves metabolic syndrome: a preliminary dietary supplementation trial. *Nutr J* 2012;11:4.