

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.jfda-online.com

Review Article

A review on the analysis of ingredients with health care effects in health food in Taiwan

Pai-Wen Wu^{*}

Food and Drug Administration, Ministry of Health and Welfare, Taipei, Taiwan

ARTICLE INFO

Article history:

Received 6 November 2014

Received in revised form

1 March 2015

Accepted 26 March 2015

Available online 23 May 2015

Keywords:

analysis

health food

ingredients with health care effects

ABSTRACT

This review article discusses the analysis of ingredients with health care effects in health food in Taiwan. The top 10 items on the list of registered health food products up to 2014 in Taiwan are described, including monocolin K, ω -3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid), β -glucans, inulin, catechins, oligosaccharides, resistant maltodextrin, amino acids, medium chain fatty acids, and polysaccharides. Some analytical methods for the analysis of ingredients with health care effects are announced to the public on the website of health food section of the Taiwan Food and Drug Administration for the application and the postmarket surveillance of health food. Each application of health food should include the appropriate analytical method for the analysis of the ingredient or specific compound that has the health care effect, for the sake of quality assurance. Self-management of each applicant is required for regulation, the reputation of its own, and social responsibility to the consumers.

Copyright © 2015, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan

LLC. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

"Health Food" is a legislative term used in Taiwan. The total number of products designated as health foods from 1999 to 2014 is 310, including 50 health foods with specification standards (Fig. 1) [1]. There are two tracks to apply to register a health food. Track 1 is the application for individual case review deliberated by the Health Food Advisory Committee, including safety assessments (Table 1), health care effects (Table 2), and quality stability. However, Track 2, for the specification standard, skips the reviews of the Health Food Advisory Committee due to the safety and health care effects of the ingredients. Up to now, red mold rice and fish oil are the

two health food specification standards [2]. The management system of health food in Taiwan, similar to that of Japan and China, belongs to premarket registration [3–6]. Applications should be deliberated by the Health Food Advisory Committee which examine safety, effective efficiency, and evaluate the stability of the ingredient with the health care effect.

The quality of ingredients with health care effect in health foods should be assessed in order to ensure the health benefits to the consumers. The ingredients with health care effects should be qualified and quantitated. Therefore, appropriate analytical methods for the ingredients with health care effects are necessary to assess the compliance with specifications. There are more than 100 items of ingredients with health care effects in the 310 registered health food products up to 2014 in

^{*} Division of Research and Analysis, Food and Drug Administration, No. 161-2, Kunyang Street, Nangang District, Taipei City 115-61, Taiwan, ROC.

E-mail address: tfdawu@fda.gov.tw.
<http://dx.doi.org/10.1016/j.jfda.2015.03.007>

1021-9498/Copyright © 2015, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).




Fig. 1 – Number of permitted health foods from 1999 to 2014.

Taiwan. In the regulation of health food, it is necessary to provide the analytical methods used to examine ingredients with health care effects for each application, so that the method may be recognized as having scientific reliability and accuracy [7]. Because of the complexity and diversity of matrices, the available and reliable analytical methods for various types of health food effects are not adequate for specific ingredient identification or composition confirmation. The appropriate analytical methods of ingredients with health care effects for each product application should be validated and verified. The quality of ingredients with health care effects in health food is an issue of great concern.

2. Ingredients with health care effects

There are 13 health care effects announced by the competent authority (Table 2). The experimental results represent the health care effects of the complete health food product rather than just a single compound or ingredient. Because of the diverse properties, there are various ingredients with health care effects in permitted health foods. However, for some cases, the effective components cannot be clearly identified. Thereby, the index compounds are presented for the quality control of the product. This excludes the probiotics group, the top 10 ingredients with health care effects: monacolin K, ω -3

Table 2 – Thirteen health care effects of the permitted health foods.

1. Modulating blood lipids
2. Improving gastrointestinal functions
3. Regulating immune system
4. Adjuvant modulating allergic constitution
5. Protecting bone health
6. Maintaining dental health
7. Modulating blood sugar
8. Protecting liver functions (chemically induced liver damage)
9. Reducing body fat formation
10. Anti-weariness
11. Postponing aging
12. Adjuvant modulating blood pressure
13. Promoting iron absorption

Table 3 – Top 10 ingredients with health claim/index compounds for quality control in permitted health foods.

Ingredient with health claim/index compound for quality control	Number of permitted health foods
Monacolin K	45
EPA and DHA	22
β -Glucan	18
Inulin	16
Catechins	13
Oligosaccharides	12
Resistant maltodextrin	9
Amino acids, medium fatty acids, polysaccharides	8

DHA = docosahexaenoic acid; EPA = eicosapentaenoic acid.

fatty acids (eicosapentaenoic acid and docosahexaenoic acid), β -glucans, inulin, catechins, oligosaccharides, resistant maltodextrin, amino acids, medium chain fatty acids, and polysaccharides (Table 3). For the sake of postmarketing surveillance of health foods, and the necessity of commonality for the application, there are some analytical methods of efficient compounds available to the public in the section on health food analysis on the Taiwan Food and Drug Administration (TFDA) website (Table 4) [8]. The availability of those

Table 1 – Categories of safety assessments.

Category	Content	Toxicity test
I	The raw materials of the products are conventionally consumed and the product is in the form of commonly processed food; or there is complete academic literature and reports on the toxicology and safety, as well as records of consumption, of the product	May be exempted under any of the circumstances listed on the left
II	The raw materials of the product are conventionally consumed and the product is not in the form of commonly processed food	Genotoxicity study, 28-d feeding toxicity study
III	The raw materials of the product are not conventionally consumed	Genotoxicity study, 90-d feeding toxicity study, teratogenicity
IV	The raw materials of the product are not conventionally consumed and contain carcinogen analogues	Genotoxicity study, 90-d feeding toxicity study, teratogenicity, carcinogenicity study, reproduction study

Table 4 – Analytical methods of the ingredients (index compounds/specific components) with health claims in health foods.

September 15, 2014

Effective index / specific compound	Scope	Method	Attribute	Equipment/principle
DHA, EPA	Fish oil	Method of test for eicosapentaenoic acid and docosahexaenoic acid in fish oils. MOHW (Ministry of Health and Welfare) Food No. 1021950329 Amended, September 6, 2013	Promulgated method	Analysis by using gas chromatography after saponification and methylation
Oleic acid, linoleic acid, linolenic acid	Food	Method of test for fatty acids in food. MOHW Food No. 1021950978 Amended, November 28, 2013	Promulgated method	Analysis by using GC after extraction of oil, saponification and methylation
Vitamin D	Capsule, tablet, powder	Method of test for vitamin D in food. October 6, 2013	Recommended method	Analysis using HPLC after extraction
HMG-CoA inhibitor (Monacolin K)	Red mold powder, cereal powder	Red mold rice health food with specification standard. DOH Food No. 0960406448 promulgated, December 24, 2007	Promulgated method	Analysis using HPLC after extraction
Citrinin	Red mold powder, cereal powder	Method of test for mycotoxin in foods – test of citrinin. MOHW Food No. 1021950329 Amended, September 6, 2013	Promulgated method	Analysis using HPLC after extraction
Catechins	Tea beverage	Method of test for catechins in drink. November 16, 2010	Recommended method	Analysis using HPLC after dilution
β-carotene	Algae powder	β-Carotene in supplements and raw materials reversed-phase HPLC method. AOAC Official Method 2005.07	Recommended method	Analysis using HPLC after extraction
Xylitol	Chewing gum	Method for test xylitol in chewing gum. June 30, 2010	Recommended method	Analysis using HPLC after extraction
Calcium	Tablet, milk powder, liquid milk	General method of test for heavy metals. MOHW Food No. 1031901169 Amended, August 25, 2014	Promulgated method	According to the conditions of element species, sample matrix, quantitative limits and laboratory equipment, a complete analysis method is combined by choosing the appropriate digestion method and content determination method.
Inulin, fructan	Beverage, powder food	Measurement of fructan in foods-enzymatic / spectrophotometric method AOAC official method 999.03	Recommended method	Analysis using spectrophotometry after enzymatic treatment
b-glucans	Cereal food	β-D-glucan in oats, streamlined enzymatic-spectrophotometric method. AOAC Official Method 995.16	Recommended method	Analysis using spectrophotometry after enzymatic treatment
Lactic acid bacteria	Dairy product, milk powder, capsule, tablet, powder, crystalline powder	Methods of test for food microorganisms-test of lactic acid bacteria. MOHW Food No. 1021950329 Amended, September 6, 2013	Promulgated method	After serial dilution, cultivated with selective medium
<i>Bifidobacterium longum</i>		Methods of test for food microorganisms - Test of lactic acid bacteria - <i>Bifidobacterium longum</i> . June 6, 2013	Recommended method	Strains identification with real-time polymerase chain reaction after isolation and purification and
<i>Bifidobacterium lactis</i>		Methods of test for food microorganisms - Test of lactic acid bacteria - <i>Bifidobacterium lactis</i> . June 6, 2013	Recommended method	DNA extraction
<i>Bifidobacterium breve</i>		Methods of test for food microorganisms - Test of lactic acid bacteria - <i>Bifidobacterium breve</i> . June 6, 2013	Recommended method	
<i>Bifidobacterium bifidum</i>		Methods of test for food microorganisms - test of lactic acid bacteria - <i>Bifidobacterium bifidum</i> . June 6, 2013	Recommended method	
<i>Enterococcus faecalis</i>		Methods of test for food microorganisms - test of lactic acid bacteria - <i>Enterococcus faecalis</i> . November 22, 2012	Recommended method	
<i>Streptococcus thermophilus</i>		Methods of test for food microorganisms - test of lactic acid bacteria - <i>Streptococcus thermophilus</i> . November 22, 2012	Recommended method	

When the sample matrix is not the same as described in the scope of the recommended method, the assessment should be evaluated according to the *Guidelines for the Validation of Chemical Methods for Food or Guidelines for Laboratory Quality Management – Test Results of Quality Control (Microbial Field)*.

If there is any interference on test result, a further study is required.

AOAC = Association of Official Analytical Chemists; DHA = docosahexaenoic acid; EPA = eicosapentaenoic acid; GC = gas chromatography; HPLC = high performance liquid chromatography.

analytical methods is based on the ingredients can be identified; applied more than two permitted applications or applicants, or for the other reasons. The analytical methods on the TFDA website are either promulgated methods for the foundations of food sanitation standards and health food with specification standards (fish oil and red mold rice); or recommended methods modified and validated as necessary, after which the limit of quantitation (LOQ) should be equal to the original references. In addition, the international published methods, such as the Association of Official Analytical Chemists (AOAC) official methods of analysis, are recognized after validation for each application. Some analytical methods for ingredients with health care effect in health foods are interpreted in detail as follows.

2.1. Monacolin K in red mold rice

As reported, monacolin K (MK) can lower blood cholesterol concentrations effectively [9,10]. There are 45 health food products that claim MK as an ingredient with a health care effect, of which 33 belong to specification standards [1]. The daily intake of MK for red mold rice health food with a specification standard should be in the range of 4.8–10 mg [11], which is summed up with its lactone form (MKL) and hydroxy acid form (MKA). The physiologically active form of MK is its hydroxyl acid form converted from the lactone ring. MKL transfers easily into MKA with bioactivity *in vivo* [12–14]. Both forms coexist in red mold rice [15,16]. The ratios of MKL over MKA are in the range of 20:1 to 3:1 for the 12 red mold rice products (data are not published). A high performance liquid chromatography-diode array detection (HPLC-DAD) method for determining monacolin K in red mold rice capsules is promulgated in 2011 [17]. The MKL acetonitrile stock standard solution is stored at -20°C , and the working standard solution and sample are treated with methanol. MKL and MKA remain at 95% and 98%, respectively, after 24 hours. Nevertheless, MK methanol working standard solution should be prepared fresh daily. Moreover, analysis of MK should be completed as soon as possible because of MKL standard acetonitrile stock solution that could be converted to MKA in the appropriate alkali condition. The MK contents of the 12 health food products with red mold rice powder are in the range 1.49–25.0 mg/g [18]. After calculation of the net weight of each capsule and daily intake dose, nine specification standard products are in the range of 5.4–10 mg, which are compliant with the regulations.

In addition, the products applied to red mold rice health food with a specification standard should not only examine MK, but also citrinin, a type of nephrotoxic mycotoxin, of which content should less than 2 ppm and analyzed by using HPLC with fluorescence detection [10,19].

2.2. Fatty acids

Functional fatty acids, including linoleic acid, linolenic acid, γ -linolenic acid, and eicosapentaenoic acid (EPA) docosahexaenoic acid (DHA), are applied as the effective ingredients/index compounds for quality control in the permitted Health Food, which claim health benefits of lowering lipid or cholesterol in blood. The oil sample or the oil extraction from foods are done using the processes of hydrolysis, methylation to fatty acid

methyl esters, and analysis using capillary gas chromatography with flame ionization detection (GC-FID) [20–22]. The analytical methods for testing for fatty acids in food [23] and EPA and DHA in fish oils are available on the TFDA website [24]. Both of these methods are catalyzed by boron trifluoride methanol solution, whereas they have different internal standards and separation conditions.

2.3. EPA ethyl ester and DHA ethyl ester in fish oil

In compliance with the regulation of standardized fish oil health food, the contents of EPA and DHA are regulated in the range of 30–50% and the daily intake allowance of them are summed up in the range of 1–2 mg/g in triacylglycerol (TG) form [25]. The size exclusion method is applied to distinguish the glyceride forms of oil [26]. EPA and DHA exist in human body in triglyceride form naturally, rather than the ethyl ester form from refined processing [27,28]. Therefore, the mild catalyzation with tetramethylammonium hydroxide (TMAH) is applied to identify and quantify EPA ethyl ester and DHA ethyl ester in fish oil [29,30].

2.4. Catechins

Catechins are a group of flavonoid compounds with some derivatives, containing biological activity related to blood lipid adjustment or reducing body fat formation [31–33]. Eleven health foods declare catechins as the effective ingredients; most of the products are tea beverages. An event occurred in 2010, where the catechin content of some tea beverages with the “Health Food” logo were noncompliant with labeling. Therefore, a HPLC method with good resolution of eight catechins (gallic catechin, epigallocatechin, catechin, epicatechin, epigallocatechin gallate, gallic catechin gallate, catechin gallate, and epicatechin gallate) and caffeine is developed and shown on the TFDA website as the recommended analytical method for application and surveillance [34].

2.5. Polysaccharides

The polysaccharides is a complicated category because of its diverse structure [35]. In permitted health food products that claim polysaccharides as the ingredients with health care effects, they usually contain ganoderma, mushroom, ginseng, or a mixture of these [7].

The analytical procedures of determination of polysaccharides usually digest these macromolecular compounds by enzymes or chemicals into smaller compounds, which are then detected using chromogenic reactions or chromatographic instruments [36–38]. However, the aforementioned methods do not seem precise enough to coordinate with the efficiency for these compounds. Recently, the analysis of the precise effective parts of polysaccharides was studied. The determination of (1,3/1,6)- β -D-glucan extracted from fungus and yeasts containing immune modulating properties [39–41], interpreted the content and the degree of branching of the branched (1,3/1,6)- β -D-glucan simultaneously without tedious isolation and purification procedures [42,43]. The requirement of effective polysaccharide compounds for

health food applications will be more precise and compliant to keep pace with developments.

In addition, there are further developments on the specific categories of effective polysaccharides described as follows.

2.6. β -Glucans

(1,3/1,4)- β -D-glucan extracted from the bran of some grains, such as oats, is a rich source of the water-soluble fiber (1,3/1,4)- β -D-glucan, which has health claims related to the effects on lowering blood cholesterol and regulating the immune system [44–46]. AOAC methods are widely adopted for the analysis of β -glucan in oats and its related products; the principle of analysis of these methods is hydrolysis by lichenase and beta-glucosidase to glucose, then measurement using a glucose oxidase–peroxidase buffer mixture [47]. The absorption response increases as the glucose amount increases. This method was found to be applicable for the measurement of β -glucan in cereal products containing high levels of glucose after pre-extraction with aqueous ethanol. However, there is no description of the definition of high glucose levels or the concentration of ethanol used. One study demonstrated that the detected amounts of β -glucan in oat products had significant differences when more than 5% glucose was added. In this situation, pretreatment with 50% ethanol resulted in steadier data. Added sucrose or milk powder in oat products did not influence the determination of β -glucan [48].

2.7. Inulin

Inulin is a heterogeneous complex of fructose polymers, and its effect is related to the colonic microflora, gastrointestinal function, or immune system regulating [49,50]. An enzymatic and spectrophotometric collocation method is recommended to measure the fructan in foods [51], as according to theory inulin is excluded while other sugars can be hydrolyzed with enzymes (sucrase, α -amylase, maltase, and pullanase) then reduced to sugar alcohol, the remaining fructan hydrolyzed by fructanase (*exo*-inulinase plus *endo*-inulinase) and measured by the *p*-hydroxybenzoic acid hydrazide (PAHBAH) collocation method for reducing sugars.

2.8. Total dietary fiber

The definitions of dietary fiber have been discussed in depth [52–54], the common definition by the American Association for Clinical Chemistry (AACC) is “dietary fiber is the edible parts of plants or analogous carbohydrates that are resistant to digestion and absorption in the human small intestine with complete or partial fermentation in the large intestine”. The analytical methods of dietary fibers depend on their scope of application, as analysis of the dietary fiber can refer to insoluble high molecular fiber [55] or soluble low molecular compounds [56–58].

For example, some food products containing resistant maltodextrin, which have health claims regarding maintenance of normal blood low density lipoprotein cholesterol concentrations or serum concentrations of triglycerides, and changes in bowel function [59], are usually analyzed with

AOAC methods, in which the total fiber is calculated as the sum of insoluble dietary fiber measured by enzymatic-gravimetric method plus the high molecular weight soluble dietary fiber and low molecular weight resistant maltodextrin measured by an HPLC method [56–58].

2.9. Oligosaccharides and sugar alcohol

Oligosaccharides such as fructooligosaccharides, isomaltooligosaccharides, and galactooligosaccharides are claimed to improve gastrointestinal function [60–62]. The analytical methods for determination of oligosaccharides include enzymatic-colorimetric and HPLC methods, and comparison with specific standards [52,63–65].

Health foods containing xylitol as a sugar replacement have health claims for maintaining dental health [66], and may be analyzed using HPLC [67].

2.10. Amino acids

Amino acids are not only the building block of proteins, but also are involved in the nutrition metabolism and oxidative defense [68,69]. Some amino acids such as alanine, arginine, glycine, isoleucine, L-arginine, leucine, taurine, and valine are listed as the effective ingredients or the quality control index compounds in the permitted health food products, the health claims of which are related to antifatigue or immunity. The analyses of amino acids are usually detected by using either HPLC-UV or an amino acid analyzer after derivatization. The other choices are to analyze directly with LC-MS/MS [70–72].

2.11. Terpenoids

Some products containing ganoderma, ginseng, fuling, *Taiwanofungus camphoratus*, or other mushrooms claim the terpenoids (ganoderic acids, ginsenosides, lanostanes, and so on) as the ingredients with health care effects in health food applications [4]. These health foods show various health benefits, such as the immunostimulating activities via induction of cytokines and enhancement of immunological effectors, and protection of liver injury induced by tetrachloromethane. Most of these specific terpenoids are analyzed using HPLC-UV or LC-MS/MS [73,74]; however, some of them appeared as possible triterpenoids by using spectrophotometry [75].

3. Conclusion

The analytical methods for the determination of the efficacy of compounds in health foods are published on the TFDA on an ongoing basis. However, they are still not adequate for all the various types of applications. According to the self-management of food businesses, the control of raw materials is as important as the analysis of final products. As well as adopting the methods for health food analysis recommended by the TFDA or recognized by international authorities, each applicant can also develop an appropriate method following the validation guidelines [76,77]. More efforts are needed to develop validated analytical methods for the consistency of labeling of these products.

Each application of a health food should be accompanied by the appropriate analytical methods used to measure its ingredients or specific compounds that have health care effects, for the sake of quality assurance. The self-management for each product of applicant is required for the regulation, the reputation of its own, and the social responsibility to the consumers.

Conflicts of interest

The author has no conflicts of interest to declare.

REFERENCES

- [1] Food and Drug Administration, Ministry of Health and Welfare. Inquiry for the permitted Health Food. Retrieved from TFDA online database. 2014. Available at: <http://consumer.fda.gov.tw/Food/InfoHealthFood.aspx?nodeID=162> [accessed 01.05.15].
- [2] Food and Drug Administration, Ministry of Health and Welfare. The Flowchart of Application for Health Food permit. 2009. Available at: <http://consumer.fda.gov.tw/Law/Detail.aspx?nodeID=518&lang=1&lawid=161> [accessed 01.05.15].
- [3] Food and Drug Administration, Ministry of Health and Welfare. Health Food control act. 2006. Available at: <http://consumer.fda.gov.tw/Law/Detail.aspx?nodeID=518&lang=1&lawid=167> [accessed 01.05.15].
- [4] Food and Drug Administration. Enforcement Rules of Health Food Control Act. 2006. Available at: <http://consumer.fda.gov.tw/Law/Detail.aspx?nodeID=518&lang=1&lawid=168> [accessed 01.05.15].
- [5] China Food and Drug Administration. Health food registration. 2014. Available at: <http://eng.sfda.gov.cn/WS03/CL0771/> [accessed 01.05.15].
- [6] Consumer affairs agency, Food labelling division. Regulatory systems of health claims in Japan. 2011. Available at: <http://www.caa.go.jp/en/pdf/syokuhin338.pdf> [accessed 01.05.15].
- [7] Ministry of Health and Welfare. Regulations for the application of Health Food permit. 2014. Available at: <https://consumer.fda.gov.tw/Law/Detail.aspx?nodeID=518&lawid=160&k=%u5065%u5EB7%u98DF%u54C1%u7533%u8A> [accessed 01.05.15].
- [8] Food and Drug Administration, Ministry of Health and Welfare. Analytical methods of the ingredients with health claim (index compounds) or specific components in health food. 2014. Available at: <http://www.fda.gov.tw/upload/133/Content/2013051511550451194.pdf> [accessed 01.05.15].
- [9] Su YH, Wang JJ, Lin TT, Pan TM. Production of the secondary metabolites γ -aminobutyric acid and monacolin K by *Monascus*. *J Ind Microbiol Biotechnol* 2003;30:41–6.
- [10] Endo A. Chemistry, biochemistry, and pharmacology of HMG-CoA reductase inhibitors. *Klin Wochenschr* 1988;66:421–7.
- [11] Food and Drug Administration, Ministry of Health and Welfare. Red mold rice health food with specification standard. 2007. Available at: <https://consumer.fda.gov.tw/Law/Detail.aspx?nodeID=518&lawid=184&k=%u7D05%u9EB4%u5065%u5EB7%u98DF%u54> [accessed 01.05.15].
- [12] Alberts AW, Chen J, Kuron G, Hunt V, Huff J, Hoffman C, Rothrock J, Lopez M, Joshua H, Harris E, Patchett A, Monaghan R, Currie S, Stapley E, Albers-Schonberg G, Hensens O, Hirshfield J, Hoofsteen K, Liesch J, Springer J. Mevinolin: a highly potent competitive inhibitor of hydroxymethylglutaryl-coenzyme a reductase and a cholesterol-lowering agent. *P Natl Acad Sci USA* 1980;77:3957–61.
- [13] Friedrich J, Zuzek M, Bencina M, Cimerman A, Strancar A, Radez I. High-performance liquid chromatographic analysis of mevinolin acid in fermentation broths. *J Chromatogr A* 1995;704:363–7.
- [14] United State Pharmacopeia and National Formulary (USP-NF). Lovastatin. Maryland, US: USP-NF; 2005.
- [15] Chairote E, Lumyong S, Chairote G. Study on cholesterol lowering compound in red yeast rice prepared from thai glutinous rice. *As J Food Ag-Ind* 2010;3:217–28. Available online at: <http://www.ajofai.info/Abstract/Study%20on%20cholesterol%20lowering%20compounds%20in%20red%20yeast%20rice%20prepared%20from%20thai%20glutinous%20rice.pdf>.
- [16] Lee CL, Wang JJ, Pang TM. Synchronous analysis method for detection of citrinin and the lactone and acid forms of monacolin K in red mold rice. *J AOAC Int* 2006;89:669–77.
- [17] Food and Drug Administration, Ministry of Health and Welfare. Method of test for monacolin k in foods. Available at: <http://www.fda.gov.tw/upload/133/Content/2013091115594568052.pdf>; 2011 [accessed 01.05.15].
- [18] Wu PW, Wu CY, Tseng SH, Chiueh LC, Shih YC. Determination on lactone form and hydroxy acid form of monacolin k in red mold rice powder. In: *Proceeding of International Conference on Food Factors*. Taipei, Taiwan; November 20–23, 2011.
- [19] Food and Drug Administration, Ministry of Health and Welfare. Method of test for mycotoxin in foods-test of citrinin. 2013. Available at: <http://www.fda.gov.tw/upload/133/Content/2013091017173721575.pdf> [accessed 01.05.15].
- [20] AOAC International. AOAC Official Method 996.06. Fat (total, saturated and unsaturated) in foods. Gaithersburg, MD, USA: AOAC; 2002.
- [21] AOAC International. Official methods of analysis 991.39. Fatty acids in encapsulated fish oils and fish oil methyl and ethyl esters. Gaithersburg, MD, USA: AOAC; 2005.
- [22] American Oil Chemists' Society (AOCS). AOCS Official Method Ce 1h–05. Determination of cis-trans-, saturated, monounsaturated and polyunsaturated fatty acids in vegetable or non-ruminant animal oils and fats by capillary GLC. Champaign, IL, USA: AOCS; 2005.
- [23] Food and Drug Administration, Ministry of Health and Welfare. Method of test for fatty acids in food. 2013. Available at: <http://www.fda.gov.tw/upload/133/Content/2013112710470954790.pdf> [accessed 01.05.15].
- [24] Food and Drug Administration, Ministry of Health and Welfare. Method of test for eicosapentaenoic acid and docosahexaenoic acid in fish oils. 2014. Available at: <http://www.fda.gov.tw/upload/133/Content/2013091017033147870.pdf> [accessed 01.05.15].
- [25] Food and Drug Administration, Ministry of Health and Welfare. Fish oil health food with specification standard. 2007. Available at: <https://consumer.fda.gov.tw/Law/Detail.aspx?nodeID=518&lawid=183> [accessed 01.05.15].
- [26] British Pharmacopoeia. Omega-3-marine triglycerides. London, GB, UK: The Stationary Office on behalf of the Medicines and Healthcare products Regulatory Agency; 2011.
- [27] Ferraro V, Cruz IB, Jorge RF, Malcata FX, Pintado ME, Castro PML. Valorisation of natural extracts from marine source focused on marine by-products: a review. *Food Res Int* 2010;43:2221–33.
- [28] Gebauer SK, Psota TL, Harris WS, Kris-Etherton PM. n-3 Fatty acid dietary recommendations and food sources to achieve essentiality and cardiovascular benefits. *Am J Clin Nutr* 2006;83:S1526–35.

- [29] Tsai CJ, Liu CC, Hung LB, Sun Pan B. TMAH-catalyzed transesterification of EPA and DHA in encapsulated fish oil products. *J Am Oil Chem Soc* 2012;89:9–16.
- [30] Wu PW, Yang TC, Tseng SH, Kao YM, Chiueh LC and Shih YC. Determination of EPA and DHA ethyl esters in fish oil. In: 3rd Asia Pacific International Conference on Food Safety, Taipei, Taiwan; October 29 – November 1, 2013.
- [31] Riemersma RA, Rice-Evans CA, Tyrrell RM, Clifford MN, Lean ME. Tea flavonoids and cardiovascular health. *QJM* 2014;94:277–82.
- [32] Higdon JV, Frei B. Tea catechins and polyphenols: health effects, metabolism, and antioxidant functions. *Crit Rev Food Sci Nutr* 2003;43:89–143.
- [33] Nagao T, Komine Y, Soga S, Meguro S, Hase T, Tanaka Y, Tokimitsu I. Ingestion of a tea rich in catechins leads to a reduction in body fat and malondialdehyde-modified LDL in men. *Am J Clin Nutr* 2005;81:122–9.
- [34] Food and Drug Administration, Ministry of Health and Welfare. Method of test for catechins in drink. 2013. Available at: <http://www.fda.gov.tw/TC/siteList.aspx?pn=12&sid=1574> [accessed 01.05.15].
- [35] Mizuno T, Sakai T, Chihara G. Health foods and medicinal usages of mushrooms. *Food Res Int* 1995;11:69–81.
- [36] Southgate DAT. Determination of carbohydrates in foods. II. Unavailable carbohydrates. *J Sci Food Agr* 1969;9:331–5.
- [37] Englyst HN. Classification and measurement of plant polysaccharides. *Anim Feed Sci Tech* 1989;23:27–42.
- [38] Căpriță R, Căpriță A. Chemical methods for the determination of soluble and insoluble non-starch polysaccharides-Review. *Anim Sci Biotechnol* 2011;44:73–80.
- [39] Volman JJ, Ramakers JD, Plat J. Dietary modulation of immune function by β -glucans. *Physiol Behav* 2008;94:276–84.
- [40] Petravic-Tominac V, Zechner-Krpan V, Grba S, Srećec S, Panjkota-Krbavcic I, Vidovic L. Biological effects of yeast β -glucan. *Agriculturae Conspectus Scientificus* 2010;75:149–58.
- [41] Jin X, Ning Y. Extraction optimization and bioactivity of polysaccharides from *Aspergillus fumigatus* AF1. *Carbohydr Polym* 2013;96:411–6.
- [42] Chang YW, Lu TJ. Molecular characterization of polysaccharides in hot-water extracts of *Ganoderma lucidum* fruiting bodies. *J Food Drug Anal* 2004;12:59–67.
- [43] Yang Y, Ye LB, Zhang JS, Liu Y, Tang Q. Structural analysis of a bioactive polysaccharide, PISP1, from the medicinal mushroom *Phellinus igniarius*. *Biosci Biotechnol Biochem* 2009;73:134–9.
- [44] Othman RA, Moghadasian MH, Jones PJ. Cholesterol-lowering effects of oat β -glucan. *Nutr Rev* 2011;69:299–309.
- [45] European Commission. On the authorisation and refusal of authorisation of certain health claims made on foods and referring to the reduction of disease risk. Commission Regulation No. 1160/2011. Retrieved May 11, 2011.
- [46] Goodridge HS, Wolf AJ, Underhill DM. β -Glucan recognition by the innate immune system. *Immunol Rev* 2009;230:38–50.
- [47] AOAC International. AOAC Official Method 995.16. β -D-Glucan in oats-stream lined enzymatic method. Washington, DC, USA: AOAC; 2005.
- [48] Wu PW, Hsieh YJ, Tseng SH, Kao YM, Chiueh LC, Shih YC. Determination of beta-glucan in oats. *Ann Rept Food Drug Res* 2013;4:136–41.
- [49] Jenkins DJA, Kendall CWC, Vuksan V. Inulin, oligofructose and intestinal function. *J Nutr* 1999;129:1431S–3S.
- [50] Watzl B, Girschbach S, Roller M. Inulin, oligofructose and immunomodulation. *Br J Nutr* 2005;93:S49–55.
- [51] AOAC International. AOAC Official Method 999.03. Measurement of fructan in foods-enzymatic/spectrophotometric method. Washington, DC, USA: AOAC; 2007.
- [52] Dietary Fiber Definition Committee. The definition of dietary fiber. *Cereal Foods World* 2001;46:112–6.
- [53] Institute of Medicine of the National Academies. Dietary reference intakes. Washington, DC: National Academy Press; 2014.
- [54] Food and Drug Administration, Ministry of Health and Welfare. Regulations on nutrition labeling for prepackaged food products. 2014. Available at: <http://www.fda.gov.tw/EN/lawContent.aspx?id=1633&chk=d4d33c7e-5eeb-40b1-9a14-b2ed1e236dc8¶m=pn%3d1%26cid%3d16%26chk%3dd49032f6-b48e-4ab3-8fb9-223dad1b0407%26pClass1%3d> [accessed 01.05.15].
- [55] AOAC International. AOAC Official Method 985.29 Total dietary fiber in foods- enzymatic gravimetric method. Gaithersburg, Maryland, USA: AOAC; 2005.
- [56] AOAC International. AOAC Official Method 2001.03 Dietary fiber containing supplemented resistant maltodextrin (RMD). Gaithersburg, Maryland, USA: AOAC; 2006.
- [57] AOAC International. AOAC Official Method 2009.01 Total dietary fiber in foods- enzymatic gravimetric liquid chromatographic method. Gaithersburg, Maryland, USA: AOAC; 2010.
- [58] AOAC International. AOAC Official Method 2011.25 Insoluble, soluble, and total dietary fiber in foods-enzymatic gravimetric liquid chromatography. Gaithersburg, Maryland, USA: AOAC; 2012.
- [59] European Food Safety Authority. Resistant maltodextrin related health claims. *EFSA Journal* 2011;9:2070–86.
- [60] European Food Safety Authority. Fructooligosaccharides (FOS) from sucrose related health claims. *EFSA Journal* 2011;9:2023–45.
- [61] European Food Safety Authority. Isomalto-oligosaccharides related health claims. *EFSA Journal* 2010;8:1801–15.
- [62] European Food Safety Authority. Galacto-oligosaccharides (GOS) related health claims. *EFSA Journal* 2011;9:2060–75.
- [63] Slegte JD. Determination of trans-galactooligosaccharides in selected food products by ion-exchange chromatography: collaborative study. *J AOAC INT* 2002;85:417–23.
- [64] Austin S, Benet T, Michaud J, Cuany D, Rohfritsch P. Determination of β -galactooligosaccharides by liquid chromatography. *Int J Anal Chem* 2014;2014. Article ID 768406, 10 pages. Available at: <http://www.hindawi.com/journals/ijac/2014/768406/> [accessed 05.25.15].
- [65] Ko J, Lee MS, Kwak BM, Ahn JH, Park JS, Kwon JH. Determination of isomaltoligosaccharides in yoghurts by using HPLC-ELSD. *Korean J Food Sci An* 2013;33:417–24.
- [66] European Food Safety Authority. Sugar replacers xylitol, sorbitol, mannitol, maltitol, lactitol, isomalt, erythritol, D-tagatose, isomaltulose, sucralose and polydextrose related health claims. *EFSA Journal* 2011;9:2076–101.
- [67] Food and Drug Administration, Ministry of Health and Welfare. Method for test xylitol in chewing gum. Available at: <http://www.fda.gov.tw/TC/siteList.aspx?pn=11&sid=1574>; 2010 [accessed 01.05.15].
- [68] Wu G. Functional amino acids in nutrition and health. *Amino Acids* 2009;45:407–11.
- [69] Li P, Yin YL, Li D, et al. Amino acids and immune function. *Br J Nutr* 2007;98:237–52.
- [70] Marfey P. Determination of D-amino acids. II. Use of a bifunctional reagent, 1,5-difluoro-2,4-dinitrobenzene. *Carlsberg Res Commun* 1984;49:591–6.
- [71] Moore S. Amino acid analysis: Aqueous dimethyl sulfoxide as solvent for the ninhydrin reaction. *J Biol Chem* 1968;243:6281–3.
- [72] Chaimbault P, Petritis K, Elfakir C, Dreux M. Determination of 20 underivatized proteinic amino acids by ion-pairing chromatography and pneumatically assisted electrospray mass spectrometry. *J Chromatogr A* 1999;855:191–202.

-
- [73] Fuzzati N. Analysis methods of ginsenosides. *J Chromatogr B* 2004;812:119–33.
- [74] Keypour S, Rafati H, Riahi H, et al. Qualitative analysis of ganoderic acids in *Ganoderma lucidum* from Iran and China by RP-HPLC and electrospray ionisation-mass spectrometry (ESI-MS). *Food Chem* 2010;119:1704–8.
- [75] Choi J, Lee KT, Ha J, Yun SY, Ko CD, Jung HJ, Park HJ. Effects of niga-ichigoside F1 and 23-hydroxytormentic acid obtained from *Rubus coreanus*. *Biol PharmBull* 2003;26:1436–41.
- [76] Food and Drug Administration, Ministry of Health and Welfare. Guidelines for the validation of chemical analytical methods for TFDA. Available at: <http://www.fda.gov.tw/TC/siteContent.aspx?sid=1861#VGMDZdoVFv0>; 2012 [accessed 01.05.15].
- [77] US Food and Drug Administration office, Department of Health and Human Service. FDA foods program guidelines for chemical methods. 2012. Available at: <http://www.fda.gov/downloads/ScienceResearch/FieldScience/UCM298730.pdf> [accessed 01.05.15].